NKMAXBio We support you, we believe in your research

Human IL-6 antibody

Catalog Number: ATGA0313

PRODUCT INFORMATION

Catalog number

ATGA0313

Clone No.

AT1F10

Product type

Monoclonal Antibody

UnitProt No.

P05231

NCBI Accession No.

NP 000591

Alternative Names

Interleukin-6, B-cell stimulatory factor 2, BSF-2, CTL differentiation factor, CDF, Hybridoma growth factor, Interferon beta-2, IFN-beta-2, IFNB2, HGF, HSF

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human Interleukin 6 (30-212aa) purified from E. coli

Isotype

IgG1 kappa

Purification Note

By protein-A affinity chromatography

Application

ELISA, WB

Usage

The antibody has been tested by ELISA, Western blot analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results. Recommended starting dilution is 1:1000.



NKMAXBIO We support you, we believe in your research

Human IL-6 antibody

Catalog Number: ATGA0313

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


Interleukin-6, or IL-6, is a multifunctional protein, 212 amino acids in length that plays critical roles in host defense, immune response and hematopoiesis. interleukin-6 (IL-6) produced by T cells, macrophages, fibroblasts, endothelial and other cells. IL-6 induces proliferation or differentiation in many cell types including B cells, thymocytes and T cells. IL-6, in concert with TGF-Beta, is important for developing Th17 responses. The IL-6 receptor is a trimeric complex composed of an IL-6-specific a chain and a homodimer of the gp130 glycoprotein common to the IL-6, IL-11, CNTF, OSM and LIF receptors. Stimulation with IL-6 leads to gp130 homodimerization and the activation of associated kinases JAK1 and JAK2. Once activated, JAK1 and JAK2 phosphorylate Stat3, causing its nuclear translocation and transcription of Stat3-responsive genes. IL-6 has also been shown to activate the Ras/MAP kinase pathway, which regulates NFIL6 transcription.

General References

Heinrich P.C., et al. (1998) Biochem J 334: 297-314 Heinrich P.C., et al. (1998) Z Ernahrungswiss 37: 43-49 Jones S.A. (2005) J Immunol 175: 3463-3468

DATA

Western blot analysis (WB)



The Human IL6 recombinant protein (10ng) was resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human Interleukin 6 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

