# NKMAXBIO We support you, we believe in your research

## **Human MEMO1 antibody**

Catalog Number: ATGA0566

## **PRODUCT INFORMATION**

## Catalog number

ATGA0566

#### Clone No.

AT1E9

## **Product type**

Monoclonal antibody

#### UnitProt No.

Q9Y316

#### **NCBI Accession No.**

NP 057039

#### **Alternative Names**

Protein MEMO1, C2orf4, CGI-27, MEMO, NS5ATP7

#### **Additional Information**

This product was produced from tissue culture supernatant.

## **PRODUCT SPECIFICATION**

### **Antibody Host**

Mouse

#### **Reacts With**

Human

#### Concentration

1mg/ml (determined by BCA assay)

#### **Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

## **Immunogen**

Recombinant human MEMO1 (1-297aa) purified from E. coli

## Isotype

IgG1 kappa

## **Purification Note**

By protein-A affinity chromatography

## **Application**

ELISA, WB, FACS

### **Usage**

The antibody has been tested by ELISA, Western blot and FACS analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



# NKMAXBio We support you, we believe in your research

## **Human MEMO1 antibody**

Catalog Number: ATGA0566

#### **Storage**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

## **BACKGROUND**

#### **Description**


MEMO1 (Mediator of ErbB2-driven cell motility 1) is a 297 amino acid protein. It is thought to relax extracellular chemotactic signals that are targeted at the microtubule cytoskeleton, thereby controlling cell migration. Additionally, MEMO1 is required for breast carcinoma migration, suggesting an important role in tumorigenesis. The MEMO1-RHOA-DIAPH1 signaling pathway plays an important role in ERBB2-dependent stabilization of microtubules at the cell cortex. It controls the localization of APC and CLASP2 to the cell membrane, via the regulation of GSK3B activity.

#### **General References**

Marone R., et al. (2004) Nat Cell Biol 6(6):515-522. Meira M., et al. (2009) J Cell Sci 122:787-797. Zaoui K., et al. (2010) Proc Natl Acad Sci USA 107(43):18517-18522.

## **DATA**

## Western blot analysis (WB)



The cell lysates(40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human MEMO1 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.


Lane 1.: Jurkat cell lysate Lane 2.: U87MG cell lysate

Flow cytometry (FACS)

## NKMAXBio We support you, we believe in your research

# **Human MEMO1 antibody**

Catalog Number: ATGA0566



Flow cytometry analysis of MEMO1 in U87MG cells. The cell was stained with ATGA0566 at 2-5ug for 1x10^6cells (red). A Goat anti mouse IgG (Alexa fluor 488) was used as the secondary antibody. Mouse monoclonal IgG was used as the isotype control (dark gray), cells without incubation with primary and secondary antibody was used as the negative control (light gray).

