NKMAXBIO We support you, we believe in your research

Recombinant mouse IL-12 p70 protein

Catalog Number: ATGP3739

PRODUCT INFORMATION

Expression system

Baculovirus

Domain

23-335aa(p40)/23-215aa(p35)

UniProt No.

P43432(p40)/P43431(p35)

NCBI Accession No.

NP_001290173(p40)/NP_032377(p35)

Alternative Names

IL12A, IL12B, CLMF p35, CLMF p40, Cytotoxic lymphocyte maturation factor 35 kDa subunit, Cytotoxic lymphocyte maturation factor 40 kDa subunit, Interleukin-12 alpha chain, Interleukin-12 beta chain, NK cell stimulatory factor chain 1, p35, NK cell stimulatory factor chain 2, p40, NKSF1, NKSF2

PRODUCT SPECIFICATION

Molecular Weight

35.7 kDa (p40, 313aa)

22.5 kDa (p35, 199aa)

The heterodimeric II12 has a predicted Molecular weight of 58.3 kDa (512aa).

Molecular weight on SDS-PAGE will appear higher

Concentration

0.5mg/ml (determined by Bradford assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

Purity

> 95% by SDS-PAGE

Endotoxin level

< 1 EU per 1ug of protein (determined by LAL method)

Biological Activity

The activity is determined by the IFN-g ELISA in a using NK-92 human natural killer cells. The ED50 for this effect is less or equal to 10ng/ml.

Tag

His-Tag

Application

SDS-PAGE, Bioactivity

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid



Recombinant mouse IL-12 p70 protein

Catalog Number: ATGP3739

repeated freezing and thawing cycles.

BACKGROUND

Description

IL12b/IL12a, also known as interleukin 12 (subunit beta/alpha), is a cytokine that can act as a growth factor for activated T and NK cells, enhance the lytic activity of NK/lymphokine-activated killer cells, and stimulate the production of IFN-gamma by resting PBMC. It is a key mediator of cellular-immunity and induces the differentiation of Th1 cells from precursor T helper cells. As it is associated with IL23A to form the IL-23 interleukin, it induces autoimmune inflammation and is responsible for autoimmune inflammatory diseases and may be important for tumorigenesis. Recombinant mouse IL12b/IL12a protein, fused to His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

Amino acid Sequence

IL12B(p40)

MWELEKDVYV VEVDWTPDAP GETVNLTCDT PEEDDITWTS DQRHGVIGSG KTLTITVKEF LDAGQYTCHK GGETLSHSHL LLHKKENGIW STEILKNFKN KTFLKCEAPN YSGRFTCSWL VQRNMDLKFN IKSSSSSPDS RAVTCGMASL SAEKVTLDQR DYEKYSVSCQ EDVTCPTAEE TLPIELALEA RQQNKYENYS TSFFIRDIIK PDPPKNLQMK PLKNSQVEVS WEYPDSWSTP HSYFSLKFFV RIQRKKEKMK ETEEGCNQKG AFLVEKTSTE VQCKGGNVCV QAQDRYYNSS CSKWACVPCR VRS IL12A(p35)


RVIPVSGPAR CLSQSRNLLK TTDDMVKTAR EKLKHYSCTA EDIDHEDITR DQTSTLKTCL PLELHKNESC LATRETSSTT RGSCLPPQKT SLMMTLCLGS IYEDLKMYQT EFQAINAALQ NHNHQQIILD KGMLVAIDEL MQSLNHNGET LRQKPPVGEA DPYRVKMKLC ILLHAFSTRV VTINRVMGYL SSA<HHHHHHH>

General References

Zhang X., et al, (2015) J Immunol. 195:1665-1675. Zamani A., et al, (2017) Immune Netw. 17:186-191.

DATA

SDS-PAGE



3ug by SDS-PAGE visualized by coomassie blue stain.


Lane 1: reducing condition
Lane 2: non-reducing condition

Biological Activity

NKMAXBio We support you, we believe in your research

Recombinant mouse IL-12 p70 protein

Catalog Number: ATGP3739



Mouse IL12 stimulates IFN-g secretion of the NK-92 human natural killer cells. The ED50 range is less or equal to 10 ng/ml.

