

Human CD44 antibody

Catalog Number: ACD0826

PRODUCT INFORMATION

Catalog number

ACD0826

Clone No.

5C10

Product type

Monoclonal Antibody

UnitProt No.

P16070

NCBI Accession No.

NP_001001390

Alternative Names

CD44 antigen, CDw44, Epican, Extracellular matrix receptor III, ECMR-III, GP90 lymphocyte homing/adhesion receptor, HUTCH-I, Heparan sulfate proteoglycan, Hermes antigen, Hyaluronate receptor, Phagocytic glycoprotein 1, PGP-1, Phagocytic glycoprotein I, PGP-I, LHR, MDU2, MDU3, MIC4

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human CD44 (21-145aa) purified from E. coli

Isotype

IgG2b kappa

Purification Note

By protein-G affinity chromatography

Application

ELISA, WB, ICC/IF, FACS

Usage

The antibody has been tested by ELISA, Western blot, ICC/IF and FACS analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.

Human CD44 antibody

Catalog Number: ACD0826

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

CD44 is a broadly distributed transmembrane glycoprotein that plays a critical role in a variety of cellular behaviors, including adhesion, migration, invasion, and survival. CD44 mediates cell-cell and cell-matrix interactions in a large part through its affinity for hyaluronan (HA), a glycosaminoglycan constituent of extracellular matrices, but also potentially through its affinity for other ligands such as osteopontin, collagens, and matrix metalloproteinases (MMPs). Variants of CD44 are expressed in tissues during development, including embryonic epithelia. known functions of CD44 are cellular adhesion (aggregation and migration), hyaluronate degradation, lymphocyte activation, lymph node homing, myelopoiesis and lymphopoiesis, angiogenesis, and release of cytokines. The functions of CD44 are principally dependant on cellular adhesion in one setting or another.

General References


Cichy, J. and Pure, E. (2003) J.Cell Biol. 161, 839-843.
Sneath RJ, Mangham DC (1998) Mol Pathol. 51(4):191-200.

DATA

Western blot analysis (WB)

Human CD44 antibody

Catalog Number: ACD0826




The recombinant protein (200ng) was resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human CD44 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system. Lane 1.: Recombinant human CD44 protein

Immunocytochemistry/Immunofluorescence (ICC/IF)


Human CD44 antibody

Catalog Number: ACD0826




ICC/IF analysis of CD44 in HeLa cells. The cell was stained with ACD0826 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

Flow cytometry (FACS)



Flow cytometry analysis of CD44 in HeLa cells. The cell was stained with ACD0826 at 2-5ug for 1×10^6 cells (red). A Goat anti mouse IgG (Alexa fluor 488) was used as the secondary antibody. Mouse monoclonal IgG was used as the isotype control (dark gray), cells without incubation with primary and secondary antibody was used as the negative control (light gray).



Flow cytometry analysis of CD44 in PBMC cells. The cell was stained with ACD0826 at 2-5ug for 1×10^6 cells (red). A Goat anti mouse IgG (Alexa fluor 488) was used as the secondary antibody. Mouse monoclonal IgG was used as the isotype control (dark gray), cells without incubation with primary and secondary antibody was used as the negative control (light gray).