NKMAXBIO We support you, we believe in your research

Human 14-3-3 epsilon antibody

Catalog Number: ATGA0124

PRODUCT INFORMATION

Catalog number

ATGA0124

Clone No.

AT4F8

Product type

Monoclonal Antibody

UnitProt No.

P62258

NCBI Accession No.

NP 006752

Alternative Names

YWHAE, tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein epsilon, 14-3-3E, HEL2, KCIP-1, MDCR, MDS

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human14-3-3 epsilon (1-255aa) purified from E. coli

Isotype

IgG2b kappa

Purification Note

By protein-A affinity chromatography

Application

ELISA, WB, ICC/IF

Usage

The antibody has been tested by ELISA, Western blot analysis, Flow cytometry and ICC/IF to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



NKMAXBIO We support you, we believe in your research

Human 14-3-3 epsilon antibody

Catalog Number: ATGA0124

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


The 14-3-3 family of proteins plays a key regulatory role in signal transduction, checkpoint control, apoptotic and nutrient-sensing pathways. 14-3-3 proteins are highly conserved and ubiquitously expressed. There are at least seven isoforms, beta, gamma, epsilon, sigma, zeta, tau and eta that have been identified in mammals. The 14-3-3 epsilon, a subtype of the 14-3-3 family of proteins, was thought to be brain and neuron-specific. It has been shown to interact with CDC25 phosphatases, RAF1 and IRS1 proteins, suggesting its role in diverse biochemical activities related to signal transduction, such as cell division and regulation of insulin sensitivity. It has also been implicated in the pathogenesis of small cell lung cancer.


General References

Oriente F., et al. (2005) J Biol Chem. 280(49): 40642-9. Conklin D., et al. (1995) PNAS. 92(17): 7892-6.

DATA

Western blot analysis (WB)





The cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human 14-3-3 epsilon antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: HeLa cell lysate Lane 2.: Jurkat cell lysate Lane 3.: 293T cell lysate Lane 4.: A549 cell lysate

Lane 5.: Mouse Brain Tissue lysate

The Recombinant Protein (50ng) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human 14-3-3 epsilon antibody (1:1000). Proteins were visualized using a goat antimouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: 14-3-3 Zeta Recombinant Protein Lane 2.: 14-3-3 Beta Recombinant Protein Lane 3.: 14-3-3 Epsilon Recombinant Protein Lane 4.: 14-3-3 Eta Recombinant Protein Lane 5.: 14-3-3 Gamma Recombinant Protein Lane 6.: 14-3-3 Tau Recombinant Protein


Immunocytochemistry/Immunofluorescence (ICC/IF)



NKMAXBio We support you, we believe in your research

Human 14-3-3 epsilon antibody

Catalog Number: ATGA0124



ICC/IF analysis of 14-3-3 epsilon in HeLa cells line, stained with DAPI (Blue) for nucleus staining and monoclonal anti-human 14-3-3 epsilon antibody (1:100) with goat anti-mouse IgG-Alexa fluor 488 conjugate (Green).

ICC/IF analysis of 14-3-3 epsilon in A549 cells line, stained with DAPI (Blue) for nucleus staining and monoclonal anti-human 14-3-3 epsilon antibody (1:100) with goat anti-mouse IgG-Alexa fluor 488 conjugate (Green).

