NKMAXBIO We support you, we believe in your research

Human NFATC1 antibody

Catalog Number: ATGA0133

PRODUCT INFORMATION

Catalog number

ATGA0133

Clone No.

AT1C3

Product type

Monoclonal Antibody

UnitProt No.

095644

NCBI Accession No.

NP 765978

Alternative Names

nuclear factor of activated T-cells, cytosolic component 1 isoform A, NF-ATC, NFATC, NFAT2

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human NFATc1 (428-716aa) purified from E. coli

Isotype

IgG2a kappa

Purification Note

By protein-G affinity chromatography

Application

ELISA, WB, ICC/IF, FACS

Usage

The antibody has been tested by ELISA, Western blot, ICC/IF and FACS to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.

Storage



NKMAXBio We support you, we believe in your research

Human NFATC1 antibody

Catalog Number: ATGA0133

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


Nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 1 (NFATc1) is a component of the nuclear factor of activated T cells DNA-binding transcription complex which consists of at least two components: a preexisting cytosolic component that translocates to the nucleus upon T cell receptor (TCR) stimulation, and an inducible nuclear component. This protein plays a main role in the inducible expression of cytokine genes in T-cells, especially in the induction of the IL-2 or IL-4 gene transcription and functions as a major molecular target for the immunosuppressive drugs such as cyclosporine A. NFATc1 is expressed in most human primary lymphocytes and mature human T- and B-cell neoplasms.

General References

Akimzhanov A, et al., (2008) Am J Pathol 172(1):215-224. Asagiri M, et al., (2005) J Exp Med 202(9):1261-1269. Rao A, et al., (1997) Annu Rev Immunol 15:707-747.

DATA

Western blot analysis (WB)




The cell lysates(40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human NFATC1 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: 293T cell lysate

Lane 3.: NFATC1 Transfected 293T cell lysate

Immunocytochemistry/Immunofluorescence (ICC/IF)



ICC/IF analysis of NFATC1 in HeLa cells. The cell was stained with ATGA0133 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).




NKMAXBio We support you, we believe in your research

Human NFATC1 antibody

Catalog Number: ATGA0133

Flow cytometry (FACS)



Flow cytometry analysis of NFATC1 in Jurkat cells. The cell was stained with ATGA0133 at 2-5ug for 1x10^6cells (red). A Goat anti mouse IgG (Alexa fluor 488) was used as the secondary antibody. Mouse monoclonal IgG was used as the isotype control (dark gray), cells without incubation with primary and secondary antibody was used as the negative control (light gray).

