## NKMAXBio We support you, we believe in your research

## **Human CRABP2 antibody**

Catalog Number: ATGA0134

## **PRODUCT INFORMATION**

### Catalog number

ATGA0134

#### Clone No.

AT2E11

## **Product type**

Monoclonal Antibody

#### UnitProt No.

P29373

#### **NCBI Accession No.**

NP 001869

#### **Alternative Names**

Cellular retinoic acid binding protein 2, RBP6, CRABP-II, Cellular retinoic acid binding protein 2, CRABP2, RBP6, CRABP-II, Cellular retinoic acid binding protein 2 Cellular retinoic acid binding protein II, CRABPII

## **PRODUCT SPECIFICATION**

### **Antibody Host**

Mouse

#### **Reacts With**

Human

#### **Concentration**

1mg/ml (determined by BCA assay)

#### **Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

## **Immunogen**

Recombinant human CRABP2 (1-138aa) purified from E. coli

## Isotype

IgG2a kappa

#### **Purification Note**

By protein-G affinity chromatography

## **Application**

ELISA, WB, ICC/IF

#### Usage

The antibody has been tested by ELISA, Western blot and ICC/IF analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



# NKMAXBio We support you, we believe in your research

## **Human CRABP2 antibody**

Catalog Number: ATGA0134

### **Storage**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

## **BACKGROUND**

### **Description**


The cellular retinoic acid-binding protein II (CRABP-II) is involved in the conversion of vitamin A into its intracellular active form retinoic acid, which regulate the genes responsible for lipid metabolism and adipocyte differentiation. CRABP2 gene is located on chromosome 1q21-23 and this region has been linked with related disorders such as familial combined hyperlipidemia (FCHL) and type 2 diabetes mellitus.

### **General References**

Astrom, A., et al (1991). J. Biol. Chem. 266 (26), 17662-17666. Gupta, A., et al (2006). Cancer Res. 66 (16), 8100-8108.


## **DATA**

## Western blot analysis (WB)



The cell lysate (40ug) was resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human CRABP2 antibody (1:250). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: MCF-7 cell lysate



The cell lysates (10ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human CRABP2 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: 293T cell lysate

Lane 2. : CRABP2 Transfected 293T cell lysate


Immunocytochemistry/Immunofluorescence (ICC/IF)



# NKMAXBio We support you, we believe in your research

# **Human CRABP2 antibody**

Catalog Number: ATGA0134



ICC/IF analysis of CRABP2 in HeLa cells. The cell was stained with ATGA0134 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

ICC/IF analysis of CRABP2 in MCF7 cells. The cell was stained with ATGA0134 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

