NKMAXBIO We support you, we believe in your research

Human FKBP52/FKBP4 antibody

Catalog Number: ATGA0142

PRODUCT INFORMATION

Catalog number

ATGA0142

Clone No.

AT4D3

Product type

Monoclonal Antibody

UnitProt No.

Q02790

NCBI Accession No.

NP 002005

Alternative Names

FK506 binding protein 4, FK506 binding protein 4 52 kDa FK506 binding protein, FKBP 4, FKBP 52, FKBP 59, HBI, Hsp 56, HSP binding immunophilin, p52, p59 protein, Peptidylprolyl cis trans isomerase, PPlase, Rotamase, T cell FK506 binding protein 59kD

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human FkBP4 (1-459aa) purified from E. coli

Isotype

IgG2b kappa

Purification Note

By protein-G affinity chromatography

Application

ELISA, WB, ICC/IF, IHC, FACS

Usage

The antibody has been tested by ELISA, Western blot, ICC/IF, FACS and IHC analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



NKMAXBio we support you, we believe in your research

Human FKBP52/FKBP4 antibody

Catalog Number: ATGA0142

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


FkBP4, also known as FkBP52, is a member of the immunophilin protein family, which play a role in immunoregulation and basic cellular processes involving protein folding and trafficking. It is a component of unactivated mammalian steroid receptor complexes and may play a role in the intracellular trafficking of hetero-oligomeric forms of the steroid hormone receptors. It interacts with interferon regulatory factor-4 and plays an important role in immunoregulatory gene expression in B and T lymphocytes.

General References

Ostrow KL, et al., (2009) Clin Cancer Res. 15(4):1184-91. Ma D, et al., (2008) J Biol Chem. 283(38):25963-70.

DATA

Western blot analysis (WB)



The cell lysates(40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human FKBP52 /FKBP4 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.


Lane 1.: HeLa cell lysate Lane 2.: K562 cell lysate Lane 3.: Jurkat cell lysate Lane 4.: MCF7 cell lysate

Immunocytochemistry/Immunofluorescence (ICC/IF)




Human FKBP52/FKBP4 antibody

Catalog Number: ATGA0142




ICC/IF analysis of FKBP52 /FKBP4 in HeLa cells. The cell was stained with ATGA0142 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

Flow cytometry (FACS)



Flow cytometry analysis of FKBP52 /FKBP4 in HeLa cell line, staining at 2-5ug for 1×10^6 cells. The secondary antibody used goat antimouse IgG Alexa fluor 488 conjugate.

Immunohistochemistry (IHC)



Seminoma (Germ cell tumor)

Paraffin embedded sections of Seminoma (Germ cell tumor) tissue were incubated with anti-human FKBP52 /FKBP4 (1:50) for 2 hours at room temperature. Antigen retrieval was performed in 0.1M sodium citrate buffer and detected using Diaminobenzidine (DAB)

