NKMAXBio We support you, we believe in your research

Human TBCA antibody

Catalog Number: ATGA0155

PRODUCT INFORMATION

Catalog number

ATGA0155

Clone No.

AT1A5

Product type

Monoclonal Antibody

UnitProt No.

075347

NCBI Accession No.

NP 004598

Alternative Names

Tubulin folding cofactor A, Tubulin folding cofactor A, chaperonin cofactor a, tubulin specific chaperone a, TBCA, Tubulin folding cofactor A CFA, Co chaperonin associated with a & b tubulin, Cofactor A, TCP1 chaperonin cofactor A, Tubulin cofactor a

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant TBCA (1-108aa) purified from E. coli

Isotype

IgG2b kappa

Purification Note

By protein-G affinity chromatography

Application

ELISA, WB, ICC/IF, FACS

Usage

The antibody has been tested by ELISA, Western blot, ICC/IF and FACS analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



NKMAXBio We support you, we believe in your research

Human TBCA antibody

Catalog Number: ATGA0155

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


Tubulin folding cofactor A, also known as TBCA, is one of four proteins (cofactors A, D, E, and C) involved in the pathway leading to correctly folded beta-tubulin from folding intermediates. Cofactors A and D are believed to play a role in capturing and stabilizing beta-tubulin in a quasi-native confirmation. This protein is essential for cell viability and its knockdown produces a decrease in the amount of soluble tubulin, modifications in microtubules and G1 cell cycle arrest.

General References


Nolasco S, et al., (2005) FEBS Lett 579(17):3515-3524. Tian G, et al., (1996) Cell 86(2):287-296.

DATA

Western blot analysis (WB)



The cell lysate of LNCap(20ug) was resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human TBCA antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.




The cell lysates(40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human TBCA antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: HeLa cell lysate Lane 2.: K562 cell lysate


Immunocytochemistry/Immunofluorescence (ICC/IF)

Human TBCA antibody

Catalog Number: ATGA0155




ICC/IF analysis of TBCA in A549 cells. The cell was stained with ATGA0155 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).



ICC/IF analysis of TBCA in HeLa cells. The cell was stained with ATGA0155 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

Flow cytometry (FACS)



Flow cytometry analysis of TBCA in HeLa cell line, staining at 2-5ug for 1×10^6 cells. The secondary antibody used goat anti-mouse IgG Alexa fluor 488 conjugate.

