PRODUCT INFORMATION

Catalog number ATGA0178

Clone No. AT4C3

Product type Monoclonal Antibody

UnitProt No. Q96SI1

NCBI Accession No. NP_076981

Alternative Names

BTB/POZ domain-containing protein KCTD15, MGC25497, MGC2628, potassium channel tetramerisation domain containing 15

PRODUCT SPECIFICATION

Antibody Host Mouse

Reacts With Human

Concentration 1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human KCTD15 (1-234) purified from E.coli

Isotype

lgG3 kappa

Purification Note By protein-G affinity chromatography

Application

ELISA,WB,ICC/IF

Usage

The antibody has been tested by ELISA, Western blot and ICC/IF analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND


Description

KCTD15 (Potassium channel tetramerisation domain containing 15, also known as BTB/POZ domain-containing protein KCTD15) is protein that in humans is encoded by the KCTD15 gene. KCTD15 is expressed at high level in brain and hypothalamus. The potassium channel KCTD15 was identified as a genetic loci associated with higher than normal body mass index (BMI) in humans along with genes such as GNPDA2, MTCH2, FTO, and TMEM18. Single nucleotide polymorphisms (SNPs) in non-diabetic and diabetic patients showed that FTO was most strongly associated with obesity while MTCH2 and GNPDA2 were still significantly associated with higher than normal BMI levels.

General References

Willer CJ, et al. (2010) Nat. Genet, 41(1):24-34. Elks CE, et al. (2010) PLoS Med, 7(5):e1000284. Strausberg RL, et al. (2002) Proc. Natl. Acad. Sci. U.S.A, 99(26):16899-903.

DATA



Western blot analysis (WB)

The cell lysate of HeLa (35ug) was resolved by SDS-PAGE, transferred to NC membrane and probed with anti-human KCTD15 (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.


The cell lysates (10ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human KCTD15 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system. Lane 1 : 293T cell lysate Lane 2 : KCTD15 Transfected 293T cell lysate

2

NKMAX

NKMAXBiO We support you, we believe in your research Human KCTD15 antibody Catalog Number: ATGA0178

Immunocytochemistry/Immunofluorescence (ICC/IF)



Immunofluorescence of human HeLa cells stained with monoclonal anti-human KCDT15 antibody (1:500) with Texas Red (Red). Nucleus was stained by Hoechst 33342 (Blue).

