

# Human Adenosine Kinase/ADK antibody

Catalog Number: ATGA0206

## PRODUCT INFORMATION

---

**Catalog number**

ATGA0206

**Clone No.**

AT4F8

**Product type**

Monoclonal Antibody

**UnitProt No.**

P55263

**NCBI Accession No.**

NP\_006712

**Alternative Names**

ADK, AK, adenosine kinase, adenosine 5'-phosphotransferase

## PRODUCT SPECIFICATION

---

**Antibody Host**

Mouse

**Reacts With**

Human

**Concentration**

1mg/ml (determined by BCA assay)

**Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

**Immunogen**

Recombinant human ADK (22-362aa) purified from E. coli

**Isotype**

IgG1 kappa

**Purification Note**

By protein-G affinity chromatography

**Application**

ELISA, WB, ICC/IF, FACS

**Usage**

The antibody has been tested by ELISA, Western blot, ICC/IF and FACS analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.

# Human Adenosine Kinase/ADK antibody

Catalog Number: ATGA0206

## Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

## BACKGROUND

### Description


Adenosine kinase (ADK) is the key regulator of adenosine metabolism. Because of the manifold receptor-dependent actions of adenosine, tight regulation of adenosine levels is crucial. The intracellular and extracellular pools of adenosine are in dynamic exchange by equilibrative and concentrative nucleoside transporters, so extracellular concentrations of adenosine are regulated by interplay of these transporters with intracellular and extracellular enzymes of adenosine metabolism. Thus, the extracellular concentration of adenosine is enhanced by inhibition of equilibrative nucleoside transporters such as S- (4-nitrobenzyl) -6-thioinosine and cannabidiol and, stimulation of extracellular ATP breakdown and inhibition of intracellular adenosine removal.

### General References

- Mathews II , et al. (1998). *Biochemistry* 1037(45):15607-20.
- Lindberg B, Klenow H, Hansen K (1967). *J. Biol. Chem.* 242 (3): 350-6.
- CAPUTTO R (1951). *J. Biol. Chem.* 189 (2): 801-14.

## DATA

### Western blot analysis (WB)




The cell lysates of HepG2 and 293T (35ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human ADK (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

### Immunocytochemistry/Immunofluorescence (ICC/IF)


## Human Adenosine Kinase/ADK antibody

Catalog Number: ATGA0206



ICC/IF analysis of ADK in A549 cells. The cell was stained with ATGA0206 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

### Flow cytometry (FACS)



Flow cytometry analysis of ADK in A549 cell line, staining at 2-5ug for  $1 \times 10^6$  cells (red line). The secondary antibody used goat anti-mouse IgG Alexa fluor 488 conjugate. Isotype control antibody was mouse IgG (black line).