# NKMAXBIO We support you, we believe in your research

## **Human ARF1 antibody**

Catalog Number: ATGA0207

## **PRODUCT INFORMATION**

## Catalog number

ATGA0207

#### Clone No.

AT1B3

## **Product type**

Monoclonal Antibody

#### UnitProt No.

P84077

#### **NCBI Accession No.**

NP 001649

### **Alternative Names**

ADP-ribosylation factor 1, ADP ribosylation factor 1

## **PRODUCT SPECIFICATION**

## **Antibody Host**

Mouse

#### **Reacts With**

Human

#### Concentration

1mg/ml (determined by BCA assay)

#### **Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

#### **Immunogen**

Recombinant human ARF1 (1-181aa) purified from E. coli

### Isotype

IgG1 kappa

## **Purification Note**

By protein-G affinity chromatography

## **Application**

ELISA, WB, ICC/IF, FACS

## Usage

The antibody has been tested by ELISA, Western blot, ICC/IF and FACS analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



# NKMAXBio We support you, we believe in your research

## **Human ARF1 antibody**

Catalog Number: ATGA0207

## **Storage**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

## **BACKGROUND**

## **Description**


ADP-ribosylation factor 1 (ARF1) is a member of the human ARF gene family. The family member is essential and ubiquitous in eukaryotes, being involved in vesicular transport and functioning as an activator of phospholipase D and cholera toxin. The functions of ARF proteins in membrane traffic and organelle integrity are intimately tied to its reversible association with membranes and specific interactions with membrane phospholipids. ARF1 is various membrane trafficking events in the ER-Golgi system and in the maintenance of organelle structure. Inactive ARF1 (ARF1-GDP) localizes in the cytoplasm, while the active form (ARF1-GTP) localizes in the membrane.

### **General References**

Amor, J. C. et al. (1994) Nature 372: 704-708. Claude, A. et al. (1999) J. Cell Biol. 146: 71-84. Derrien, V. et al. (2002) J. Cell Sci. 115: 2867-2879.

#### **DATA**


## Western blot analysis (WB)



The cell lysates and tissue lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human ARF1 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: HeLa cell lysate Lane 2.: HepG2 cell lysate Lane 3.: Brain tissue lysate Lane 4.: MCF7 cell lysate

## Immunocytochemistry/Immunofluorescence (ICC/IF)




ICC/IF analysis of ARF1 in MCF7 cells. The cell was stained with ATGA0207 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).




## **Human ARF1 antibody**

Catalog Number: ATGA0207



ICC/IF analysis of ARF1 in U87MG cells. The cell was stained with ATGA0207 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

## Flow cytometry (FACS)



Flow cytometry analysis of ARF1 in MCF7 cell line, staining at 2-5ug for 1x10^6cells (red line). The secondary antibody used goat antimouse IgG Alexa fluor 488 conjugate. Isotype control antibody was mouse IgG (black line).