## NKMAXBio We support you, we believe in your research

# **Human EGFR antibody**

Catalog Number: ATGA0212

## **PRODUCT INFORMATION**

## Catalog number

ATGA0212

#### Clone No.

AT2H8

## **Product type**

Monoclonal Antibody

#### UnitProt No.

P00533

#### **NCBI Accession No.**

NP 005219

#### **Alternative Names**

Epidermal growth factor receptor, ERBB, ERBB1, HER1, NISBD2, PIG61, mENA, ERRP, Proto-oncogene c-ErbB-1, Receptor tyrosine-protein kinase erbB-1, erythroblastic leukemia viral (v-erb-b) oncogene homolog (avian)

## **PRODUCT SPECIFICATION**

## **Antibody Host**

Mouse

#### **Reacts With**

Human

#### **Concentration**

1mg/ml (determined by BCA assay)

#### **Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

## **Immunogen**

Recombinant human EGFR (424-605aa) purified from E. coli

## Isotype

IgG2b kappa

### **Purification Note**

By protein-G affinity chromatography

## **Application**

ELISA, WB, ICC/IF

#### Usage

The antibody has been tested by ELISA, Western blot and ICC/IF analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



# NKMAXBio We support you, we believe in your research

# **Human EGFR antibody**

Catalog Number: ATGA0212

## **Storage**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

## **BACKGROUND**

## **Description**


Epidermal growth factor receptor (EGFR) is a 170 kDa membrane protein first identified as a binding partner of EGF. Also, EGFR phosphorylated itself (autophosphorylation) in response to EGF treatment. The overexpression of EGFR in various epithelial tumors was widely reported and it has been substantiated that deregulation of EGFR itself and its signaling pathways an important role in human cancers.

## **General References**

Cohen S, (1964) Nat Cancer Inst Monogr, 13: 13-27. Toyoda H, et al. (1995) J Biol Chem, 270: 7495-500. Carpenter G, et al. (1978) Nature, 276: 409-10.


## **DATA**

## Western blot analysis (WB)



The cell lysates of HaCaT were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human EGFR (1:250). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

## Immunocytochemistry/Immunofluorescence (ICC/IF)



Immunofluorescence of HaCaT cells stained with Hoechst 3342 (Blue) or phalloidin-TRITC for nucleus or actin staining, respectably. monoclonal anti-human EGFR antibody (1:250) was stained with alexa 488 (green).

