NKMAXBIO We support you, we believe in your research

Human PIG3/TP53I3 antibody

Catalog Number: ATGA0227

PRODUCT INFORMATION

Catalog number

ATGA0227

Clone No.

AT1C9

Product type

Monoclonal Antibody

UnitProt No.

Q53FA7

NCBI Accession No.

NP 671713

Alternative Names

Quinone oxidoreductase PIG3, NADPH:quinone reductase PIG3, Tumor protein p53-inducible protein 3, p53-induced gene 3 protein

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human TP53I3 (1-332aa) purified from E. coli

Isotype

IgG2a kappa

Purification Note

By protein-G affinity chromatography

Application

ELISA, WB, ICC/IF

Usage

The antibody has been tested by ELISA, Western blot and ICC/IF analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



NKMAXBIO We support you, we believe in your research

Human PIG3/TP53I3 antibody

Catalog Number: ATGA0227

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


TP53I3 participates in the generation of reactive oxygen species (ROS). TP53I3 has low NADPH-dependent naphtoquinone reductase activity, with a preference for 1, 2-naphtoquinone over 1, 4-naphtoquinone. TP53I3 has low NADPH-dependent diamine reductase activity (in vitro). TP53I3 is localized to the cytoplasm and induced in primary, non-transformed and transformed cell cultures after exposure to genotoxic agents. TP53I3 microsatellite polymorphism is associated with differential susceptibility to cancer.

General References

Porte S, et al. (2009) J Biol Chem.. 284(25):17194-205. Flatt PM. et al. (2000) Cancer Lett. 156(1):63-72. J Biol Chem. (2009 Jun 19) 284(25):17194-205.


DATA

Western blot analysis (WB)



The Cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human TP53I3 antibody (1:3000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1. : A549 cell lysate Lane 2. : SW480 cell lysate



The Cell lysates (1ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human TP53I3 antibody (1:3000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: 293T cell lysate

Lane 2.: TP53I3 Transfected 293T cell lysate


Immunocytochemistry/Immunofluorescence (ICC/IF)



NKMAXBio We support you, we believe in your research

Human PIG3/TP53I3 antibody

Catalog Number: ATGA0227



ICC/IF analysis of TP53I3 in A549 cells. The cell was stained with ATGA0227 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

