

# Human PIG3/TP53I3 antibody

Catalog Number: ATGA0227

## PRODUCT INFORMATION

---

**Catalog number**

ATGA0227

**Clone No.**

AT1C9

**Product type**

Monoclonal Antibody

**UnitProt No.**

Q53FA7

**NCBI Accession No.**

NP\_671713

**Alternative Names**

Quinone oxidoreductase PIG3, NADPH:quinone reductase PIG3, Tumor protein p53-inducible protein 3, p53-induced gene 3 protein

## PRODUCT SPECIFICATION

---

**Antibody Host**

Mouse

**Reacts With**

Human

**Concentration**

1mg/ml (determined by BCA assay)

**Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

**Immunogen**

Recombinant human TP53I3 (1-332aa) purified from E. coli

**Isotype**

IgG2a kappa

**Purification Note**

By protein-G affinity chromatography

**Application**

ELISA, WB, ICC/IF

**Usage**

The antibody has been tested by ELISA, Western blot and ICC/IF analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.

# Human PIG3/TP53I3 antibody

Catalog Number: ATGA0227

## Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

## BACKGROUND

### Description

TP53I3 participates in the generation of reactive oxygen species (ROS). TP53I3 has low NADPH-dependent naphtoquinone reductase activity, with a preference for 1, 2-naphtoquinone over 1, 4-naphtoquinone. TP53I3 has low NADPH-dependent diamine reductase activity (in vitro). TP53I3 is localized to the cytoplasm and induced in primary, non-transformed and transformed cell cultures after exposure to genotoxic agents. TP53I3 microsatellite polymorphism is associated with differential susceptibility to cancer.

### General References


Porte S, et al. (2009) J Biol Chem.. 284(25):17194-205.

Flatt PM. et al. (2000) Cancer Lett. 156(1):63-72.

J Biol Chem. (2009 Jun 19) 284(25):17194-205.

## DATA


### Western blot analysis (WB)



The Cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human TP53I3 antibody (1:3000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1. : A549 cell lysate

Lane 2. : SW480 cell lysate



The Cell lysates (1ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human TP53I3 antibody (1:3000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.


Lane 1.: 293T cell lysate

Lane 2.: TP53I3 Transfected 293T cell lysate

### Immunocytochemistry/Immunofluorescence (ICC/IF)

## Human PIG3/TP53I3 antibody

Catalog Number: ATGA0227



ICC/IF analysis of TP53I3 in A549 cells. The cell was stained with ATGA0227 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).