

Human LRPAP antibody

Catalog Number: ATGA0321

PRODUCT INFORMATION

Catalog number

ATGA0321

Clone No.

AT8G8

Product type

Monoclonal Antibody

UnitProt No.

P30533

NCBI Accession No.

NP_002328

Alternative Names

alpha-2-macroglobulin receptor-associated protein, A2MRAP, A2RAP, HBP44, MGC138272, MRAP, RAP

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human LRPAP1 (35-357aa) purified from E. coli

Isotype

IgG1 kappa

Purification Note

By protein-A affinity chromatography

Application

ELISA, WB, ICC/IF

Usage

The antibody has been tested by ELISA, Western blot and ICC/IF analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.

Storage

Human LRPAP antibody

Catalog Number: ATGA0321

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


Alpha-2-macroglobulin receptor-associated protein, also known as low density lipoprotein receptor-related protein-associated protein 1, RAP and LRPAP1, is a 39 kDa protein and a member of the alpha-2-MRAP family. LRPAP1 is a receptor antagonist that interacts with several members of the low density lipoprotein (LDL) receptor gene family. Upon binding to these receptors, LRPAP1 inhibits all ligand interactions with the receptors. LRPAP1 is present on cell surface forming a complex with the alpha-2-macroglobulin receptor heavy and light chains. LRPAP1 is present on the cell surface, where it is an effective inhibitor of ligand binding to the kidney glycoprotein 330 (gp330) yet it is most abundant in the endoplasmic reticulum lumen and may function in receptor folding and/or trafficking.

General References

Kounnas. M.Z., et al. (1992) J Biol Chem 267(29):21162-21166.
Bu. G., et al. (1995) EMBO J 14(10):2269-2280.


DATA

Western blot analysis (WB)



The tissue lysate (40ug) was resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human LRPAP antibody (1:500). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.
Lane 1.: Mouse kidney tissue lysate

Immunocytochemistry/Immunofluorescence (ICC/IF)



ICC/IF analysis of LRPAP in HeLa cells. The cell was stained with ATGA0321 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).