

# Human NPM1 antibody

Catalog Number: ATGA0345

## PRODUCT INFORMATION

---

**Catalog number**

ATGA0345

**Clone No.**

AT23F1

**Product type**

Monoclonal Antibody

**UnitProt No.**

P06748

**NCBI Accession No.**

NP\_002511

**Alternative Names**

Nucleophosmin, B23, NPM

## PRODUCT SPECIFICATION

---

**Antibody Host**

Mouse

**Reacts With**

Human

**Concentration**

1mg/ml (determined by BCA assay)

**Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

**Immunogen**

Recombinant human NPM1(1-294aa) purified from E. coli

**Isotype**

IgG1 kappa

**Purification Note**

By protein-A affinity chromatography

**Application**

ELISA, WB, ICC/IF, FACS

**Usage**

The antibody has been tested by ELISA, Western blot, ICC/IF and FACS analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.

# Human NPM1 antibody

Catalog Number: ATGA0345

## Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

## BACKGROUND

### Description

NPM1 is associated with nucleolar ribonucleoprotein structures and bind single-stranded and double-stranded nucleic acids, but it binds preferentially G-Quadruplex forming nucleic acids. It is involved in the biogenesis of ribosomes and may assist small basic proteins in their transport to the nucleolus. Its regulation through SUMOylation (by SENP3 and SENP5) is another facet of the proteins's regulation and cellular functions. It is located in the nucleolus, but it can be translocated to the nucleoplasm in case of serum starvation or treatment with anticancer drugs. The protein is phosphorylated. Chromosomal aberrations involving NPM1 were found in patients with non-Hodgkin lymphoma, acute promyelocytic leukemia, myelodysplastic syndrome, and acute myelogenous leukemia. It has been found in the cytoplasm in patients with primary acute myelogenous leukemia.


### General References

Maiguel D.A., et al.(2004) Mol cell Biol. 24: 3703-3711.

Colombo E., et al. (2005) Mol cell Biol. 25: 8874-8886.

## DATA

### Western blot analysis (WB)



The recombinant protein (50ng), cell lysates(40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human NPM1 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: Recombinant human NPM1 protein


Lane 2.: Jurkat cell lysate

Lane 3.: 293T cell lysate

Lane 4.: HeLa cell lysate

Lane 5.: HepG2 cell lysate

### Immunocytochemistry/Immunofluorescence (ICC/IF)




ICC/IF analysis of NPM1 in WiDr cells. The cell was stained with ATGA0345 (1:200). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

### Flow cytometry (FACS)

## Human NPM1 antibody

Catalog Number: ATGA0345



Flow cytometry analysis of NPM1 in WiDr cells. The cell was stained with ATGA0345 at 2-5ug for  $1 \times 10^6$  cells (red). A Goat anti mouse IgG (Alexa fluor 488) was used as the secondary antibody. Mouse monoclonal IgG was used as the isotype control (blue), cells without incubation with primary and secondary antibody was used as the negative control (light gray).