NKMAXBio We support you, we believe in your research

Human GOT2 antibody

Catalog Number: ATGA0357

PRODUCT INFORMATION

Catalog number

ATGA0357

Clone No.

AT8B11

Product type

Monoclonal Antibody

UnitProt No.

P00505

NCBI Accession No.

NP 002071

Alternative Names

Aspartate aminotransferase 2 mitochondrial, Aspartate aminotransferase 2, mitochondrial, KAT4, KATIV, mitAAT

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human GOT2 (30-430aa) purified from E.coli

Isotype

IgG3 kappa

Purification Note

By protein-A affinity chromatography

Application

ELISA, WB, ICC/IF, FACS

Usage

The antibody has been tested by ELISA, Western blot, ICC/IF and FACS analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



NKMAXBIO We support you, we believe in your research

Human GOT2 antibody

Catalog Number: ATGA0357

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


Glutamic-oxaloacetic transaminase is a pyridoxal phosphate-dependent enzyme which exists in cytoplasmic and inner-membrane mitochondrial forms, GOT1 and GOT2, respectively. GOT plays a role in amino acid metabolism and the urea and tricarboxylic acid cycles. Also, GOT2 is a major participant in the malate-aspartate shuttle, which is a passage from the cytosol to the mitochondria. The two enzymes are homodimeric and show close homology. GOT2 has been seen to have a role in cell proliferation, especially in terms of tumor growth.

General References

Yang H., et al. (2015) The EMBO journal. 34(8): 1110-1125. Guidetti P., et al. (2007) Journal of neurochemistry. 102(1): 103-111.

DATA

Western blot analysis (WB)



The recombinant human GOT2 (50ng), cell and tissue lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human GOT2 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: GOT2 Recombinant protein

Lane 2.: Ramos cell lysate Lane 3.: 293T cell lysate Lane 4.: Liver tissue lysate


The cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human GOT2 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: HeLa cell lysate Lane 2.: HepG2 cell lysate Lane 3.: Balb/3T3 cell lysate



Human GOT2 antibody


Catalog Number: ATGA0357



The recombinant proteins (20ng) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human GOT2 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.


Lane 1.: Recombinant Human GOT1 Lane 2.: Recombinant Human GOT2

Immunocytochemistry/Immunofluorescence (ICC/IF)



ICC/IF analysis of GOT2 in HeLa cells. The cell was stained with ATGA0357 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

Flow cytometry (FACS)



Flow cytometry analysis of GOT2 in HepG2 cell line, staining at 2-5ug for 1x10^6cells (red line). The secondary antibody used goat antimouse IgG Alexa fluor 488 conjugate. Isotype control antibody was mouse IgG (black line).

