# NKMAXBIO We support you, we believe in your research

# **Human FUBP1 antibody**

Catalog Number: ATGA0369

# **PRODUCT INFORMATION**

## Catalog number

ATGA0369

#### Clone No.

AT14F5

# **Product type**

Monoclonal Antibody

#### UnitProt No.

Q96AE4

#### **NCBI Accession No.**

NP 003893

#### **Alternative Names**

Far upstream element (FuSE) binding protein 1, FBP, FuBP

# **PRODUCT SPECIFICATION**

## **Antibody Host**

Mouse

#### **Reacts With**

Human

#### Concentration

1mg/ml (determined by BCA assay)

#### **Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

#### **Immunogen**

Recombinant human FUBP1 (279-448aa) purified from E.coli

#### Isotype

IgG2b kappa

## **Purification Note**

By protein-A affinity chromatography

# **Application**

ELISA, WB, ICC/IF

#### Usage

The antibody has been tested by ELISA, Western blot and ICC/IF analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.

#### **Storage**



# NKMAXBio We support you, we believe in your research

# **Human FUBP1 antibody**

Catalog Number: ATGA0369

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

# **BACKGROUND**

## **Description**


FUBP1 is a transcriptional regulator and fulfills an important function in the precise control of c-myc transcription. The c-Myc protein is a transcription factor which regulates the transcription of many different target genes that play a role in proliferation, cell cycle progression, differentiation, apoptosis and cell metabolism. Consequently, FUBP1 is also involved in the regulation of proliferation and differentiation, as confirmed by different experimental approaches. Knockdown of FUBP1 or expression of a dominant-negative FUBP1 (DNA-binding domain lacking effector activity) led to proliferation arrest in U2OS and Saos-2 osteosarcoma cells due to reduced c-myc expression.

#### **General References**

Duncan R., et al. (1994) Genes Dev. 8(4): 465-480. Liu J., et al. (2006) EMBO J. 25(10): 2119-2130.

## **DATA**


# Western blot analysis (WB)



The Cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human FUBP1 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: HeLa cell lysate Lane 2.: HepG2 cell lysate Lane 3.: Jurkat cell lysate

# Immunocytochemistry/Immunofluorescence (ICC/IF)



ICC/IF analysis of FUBP1 in HeLa cells. The cell was stained with ATGA0369 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

