NKMAXBio We support you, we believe in your research

Human COX5A antibody

Catalog Number: ATGA0382

PRODUCT INFORMATION

Catalog number

ATGA0382

Clone No.

AT7A6

Product type

Monoclonal Antibody

UnitProt No.

P20674

NCBI Accession No.

NP 004246

Alternative Names

cytochrome c oxidase subunit Va, cytochrome c oxidase subunit Va, COX, COX-VA, VA

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human COX5A (42-150aa) purified from E. coli

Isotype

IgG1 kappa

Purification Note

By protein-A affinity chromatography

Application

ELISA, WB, ICC/IF

Usage

The antibody has been tested by ELISA, Western blot analysis, ICC/IF and Flow cytometry to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



NKMAXBIO We support you, we believe in your research

Human COX5A antibody

Catalog Number: ATGA0382

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


COX5A, also known as COX, COX-VA and VA, belongs to the cytochrome c oxidase subunit 5A family. Cytochrome c oxidase (COX), the terminal component of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. This component is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, and the nuclear-encoded subunits may function in the regulation and assembly of the complex. COX5A is the heme A-containing chain of cytochrome c oxidase, the terminal oxidase in mitochondrial electron transport.

General References

Lenka N., et al. (1998) Prog Nucleic Acid Res Mol Biol. 61: 309-344. Schmidt TR., et al. (2002) Gene. 286: 13-19.

DATA


Western blot analysis (WB)



The Cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human COX5A antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1. : A431 cell lysate Lane 2. : HeLa cell lysate

Immunocytochemistry/Immunofluorescence (ICC/IF)



ICC/IF analysis of COX5A in HeLa cells line, stained with DAPI (Blue) for nucleus staining and monoclonal anti-human COX5A antibody (1:100) with goat anti-mouse IgG-Alexa fluor 488 conjugate (Green).

