

Human GPI antibody

Catalog Number: ATGA0390

PRODUCT INFORMATION

Catalog number

ATGA0390

Clone No.

AT22G2

Product type

Monoclonal Antibody

UnitProt No.

P06744

NCBI Accession No.

NP_000166

Alternative Names

Glucose-6-phosphate isomerase, GPI, AMF, GNPI, NLK, PGI, PHI, SA36, EC 5.3.1.9, Neuroleukin, Phosphoglucose isomerase, Phosphohexose isomerase, SA 36, Sperm antigen 36, G6PI, Glucose-6-phosphate isomerase

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human GPI (1-558aa) purified from E. coli

Isotype

IgG2a kappa

Purification Note

By protein-A affinity chromatography

Application

ELISA, WB, FACS, ICC/IF

Usage

The antibody has been tested by ELISA, Western blot analysis, Flow cytometry and ICC/IF to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.

Human GPI antibody

Catalog Number: ATGA0390

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

Glucose-6-phosphate isomerase (GPI), also known as phosphoglucose isomerase (PGI) or phosphohexose isomerase (PHI), is an enzyme. In the cytoplasm, it functions as a glycolytic enzyme (glucose-6-phosphate isomerase) that interconverts glucose-6-phosphate (G6P) and fructose-6-phosphate (F6P). Extracellularly, it functions as a neurotrophic factor that promotes survival of skeletal motor neurons and sensory neurons, and as a lymphokine that induces immunoglobulin secretion. GPI is also referred to as autocrine motility factor (AMF) based on an additional function as a tumor-secreted cytokine and angiogenic factor. Defects in GPI are the cause of nonspherocytic hemolytic anemia, and a severe enzyme deficiency can be associated with hydrops fetalis, immediate neonatal death and neurological impairment. Alternative splicing results in multiple transcript variants.


General References

Cordeiro AT., et al. (2003) *Biochimica et biophysica acta*. 1645(2): 117-122.

Kugler W., et al. (2000) *Baillieres Best Pract Res Clin Haematol*. 13(1): 89-101.

DATA

Western blot analysis (WB)



The Cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human GPI antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1. : HeLa cell lysate

Lane 2. : HepG2 cell lysate

Lane 3. : A549 cell lysate


Lane 4. : Jurkat cell lysate

Lane 5. : U87MG cell lysate

Lane 6. : Raji cell lysate

Lane 7. : MCF7 cell lysate

Immunocytochemistry/Immunofluorescence (ICC/IF)




ICC/IF analysis of GPI in HeLa cells line, stained with DAPI (Blue) for nucleus staining and monoclonal anti-human GPI antibody (1:100) with goat anti-mouse IgG-Alexa fluor 488 conjugate (Green).

Flow cytometry (FACS)

Human GPI antibody

Catalog Number: ATGA0390



Flow cytometry analysis of GPI in HeLa cell line, staining at 2-5ug for 1x10⁶ cells (red line). The secondary antibody used goat anti-mouse IgG Alexa fluor 488 conjugate. Isotype control antibody was mouse IgG (black line).