NKMAXBio We support you, we believe in your research

Human SGTA antibody

Catalog Number: ATGA0408

PRODUCT INFORMATION

Catalog number

ATGA0408

Clone No.

AT19E8

Product type

Monoclonal Antibody

UnitProt No.

043765

NCBI Accession No.

NP 003012

Alternative Names

Small glutamine-rich tetratricopeptide repeat-containing protein alpha, alphaSGT, hSGT, SGT, Small glutamine-rich tetratricopeptide repeat-containing protein alpha Small glutamine rich tetratricopeptide repeat TPR containing alpha, uBP, Vpu binding protein

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human SGTA (1-313aa) purified from E. coli

Isotype

IgG1 kappa

Purification Note

By protein-A affinity chromatography

Application

ELISA, WB, ICC/IF

Usage

The antibody has been tested by ELISA, Western blot and ICC/IF analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



NKMAXBIO We support you, we believe in your research

Human SGTA antibody

Catalog Number: ATGA0408

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


SGTA, also known as SGT or hSGT, is a ubiquitously expressed protein that contains three TPR protein-protein interaction repeats. This protein is believed to function as a component of the androgen receptor (AR) - chaperone-cochaperone complex, acting as a cochaperone involved in androgen signaling. In addition, SGTA functions as a cochaperone that binds directly to HSC70 and HSP70 and regulates their ATPase activity. On the basis of its role in apoptosis and androgen signaling, SGTA is a potential candidate for PCOS (polycystic ovary syndrome), a disorder characterized by androgen excess, obesity and menstrual disturbances.

General References

Andreyeva A., et al. (2010) PLoS One. 5: e12018. Cziepluch C., et al. (1998) J Virol. 72(5): 4149-56.

DATA


Western blot analysis (WB)



The cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human SGTA antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: HeLa cell lysate Lane 2.: K562 cell lysate Lane 3.: A549 cell lysate Lane 4.: MCF7 cell lysate Lane 5.: LNCaP cell lysate Lane 6.: 293T cell lysate

Immunocytochemistry/Immunofluorescence (ICC/IF)



ICC/IF analysis of SGTA in Hep3B cells. The cell was stained with ATGA0408 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

