# NKMAXBio We support you, we believe in your research

# Human 6-phosphogluconate Dehydrogenase/PGD antibody

Catalog Number: ATGA0419

## **PRODUCT INFORMATION**

## Catalog number

ATGA0419

#### Clone No.

AT46B3

## **Product type**

Monoclonal Antibody

#### UnitProt No.

P52209

#### **NCBI Accession No.**

NP 002622

#### **Alternative Names**

6-phosphogluconate dehydrogenase, Decarboxylating, PGDH, Phosphogluconate dehydrogenase, 6PGD

# **PRODUCT SPECIFICATION**

## **Antibody Host**

Mouse

#### **Reacts With**

Human

#### Concentration

1mg/ml (determined by BCA assay)

#### **Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

#### **Immunogen**

Recombinant human PGD (1-483aa) purified from E. coli

#### Isotype

IgG2b kappa

## **Purification Note**

By protein-A affinity chromatography

# **Application**

ELISA, WB, ICC/IF

### Usage

The antibody has been tested by ELISA, Western blot analysis, Flow cytometry and ICC/IF to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



# Human 6-phosphogluconate Dehydrogenase/PGD antibody

Catalog Number: ATGA0419

### **Storage**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

# **BACKGROUND**

## **Description**


PGD (Phosphogluconate dehydrogenas), also known as 6PGD, is a 483 amino acid enzyme that is involved in the pentose phosphate shunt. Pentose is required for nucleic acid biosynthesis and the pentose phosphate cycle is a major source of NADPH. PGD deficiency increases the level of erythrocyte pyruvate kinase (PK) activity and reduces glutathione synthetase (GSH), resulting in hemolysis. Defects in PGD are generally asymptomatic and are inherited in an autosomal dominant fashion.

#### **General References**

Le TD., et al. (2010) Proc Natl Acad Sci U S A. 107(7): 3198-203. Tagen M., et al. (2009) J Immunol. 183(10): 6313-9

## **DATA**


# Western blot analysis (WB)



The cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human PGD antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: HeLa cell lysate Lane 2.: MCF7 cell lysate Lane 3.: 293T cell lysate Lane 4.: A549 cell lysate Lane 5.: Jurkat cell lysate

# Immunocytochemistry/Immunofluorescence (ICC/IF)



ICC/IF analysis of 6-phosphogluconate dehydrogenase/PGD1 in Hep3B cells. The cell was stained with ATGA0419 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

