NKMAXBio We support you, we believe in your research

Human ACAA1 antibody

Catalog Number: ATGA0429

PRODUCT INFORMATION

Catalog number

ATGA0429

Clone No.

AT9E5

Product type

Monoclonal Antibody

UnitProt No.

P09110

NCBI Accession No.

NP 001598

Alternative Names

3-ketoacyl-CoA thiolase peroxisomal, ACAA, PTHIO, THIO, 3-ketoacyl-CoA thiolase, peroxisomal

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human ACAA1 (27-424aa) purified from E. coli

Isotype

IgG2b kappa

Purification Note

By protein-A affinity chromatography

Application

ELISA, WB, ICC/IF

Usage

The antibody has been tested by ELISA, Western blot analysis, Flow cytometry and ICC/IF to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



NKMAXBIO We support you, we believe in your research

Human ACAA1 antibody

Catalog Number: ATGA0429

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


ACAA1 is a member of the thiolase family of enzymes and is involved in lipid metabolism. This protein is localized to the peroxisome and catalyzes the conversion of acyl-CoA and acetyl-CoA to 3-oxoacyl-CoA in the fatty acid oxidation pathway. ACAA1 shows high enzymatic activity in liver, kidney, intestine and white adipose tissue in rats. Deficiency of this enzyme leads to pseudo-Zellweger syndrome.

General References

Bour A., et al.(1991) Biochim Biophys Acta. 1090(1): 43-51. Patel S., et al. (2003) Eur Respir J. 22(5): 755-60.

DATA


Western blot analysis (WB)



The cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human ACAA1 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1 : HepG2 cell lysate Lane 2 : LnCap cell lysate

Immunocytochemistry/Immunofluorescence (ICC/IF)



ICC/IF analysis of ACAA1 in HeLa cells line, stained with DAPI (Blue) for nucleus staining and monoclonal anti-human ACAA1 antibody (1:100) with goat anti-mouse IgG-Alexa fluor 488 conjugate (Green).

