NKMAXBio We support you, we believe in your research

Human DECR1 antibody

Catalog Number: ATGA0431

PRODUCT INFORMATION

Catalog number

ATGA0431

Clone No.

AT3B2

Product type

Monoclonal Antibody

UnitProt No.

Q16698

NCBI Accession No.

NP 001350

Alternative Names

24-dienoyl-CoA reductase mitochondrial, 2,4-dienoyl-CoA reductase, mitochondrial, DECR, NADPH, SDR18C1

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human DECR1 (35-335aa) purified from E. coli

Isotype

IgG2b kappa

Purification Note

By protein-A affinity chromatography

Application

ELISA, WB, FACS, ICC/IF

Usage

The antibody has been tested by ELISA, Western blot analysis, Flow cytometry and ICC/IF to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



NKMAXBio We support you, we believe in your research

Human DECR1 antibody

Catalog Number: ATGA0431

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


DECR1 is a mitochondrial protein that exists as a homotetramer and belongs to the family of short-chain dehydrogenases/reductases. It functions as an auxiliary enzyme of beta-oxidation andt participates in the metabolism of unsaturated fatty enoyl-CoA esters. Specifically, DECR1 uses NADP+ to catalyze the reduction of 2, 4-dienoyl-CoA to yield trans-3-enoyl-CoA, which can then be used as an intermediate in the Krebs cycle. Additionally, DECR1 is thought to function as a tumor suppressor, possibly downregulating the expression of Neu and slowing the rate of tumorigenesis.

General References

Koivuranta KT., et al. (1994) Biochem J. 304(3): 787-92. Roe CR., et al. (1990) J Clin Invest. 85(5): 1703-7.

DATA

Western blot analysis (WB)



The cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human DECR1 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.


Lane 1.: PC3 cell lysate Lane 2.: HepG2 cell lysate

Immunocytochemistry/Immunofluorescence (ICC/IF)




Human DECR1 antibody

Catalog Number: ATGA0431



ICC/IF analysis of DECR1 in Hep3B cells line, stained with DAPI (Blue) for nucleus staining and monoclonal anti-human DECR1 antibody (1:100) with goat anti-mouse IgG-Alexa fluor 488 conjugate (Green).

Flow cytometry (FACS)



Flow cytometry analysis of DECR1 in Hep3B cell line, staining at 2-5ug for 1x106cells (red line). The secondary antibody used goat antimouse IgG Alexa fluor 488 conjugate. Isotype control antibody was mouse IgG (black line).

