

# Human Cyclophilin 40/PPID antibody

Catalog Number: ATGA0473

## PRODUCT INFORMATION

---

**Catalog number**

ATGA0473

**Clone No.**

AT1B8

**Product type**

Monoclonal Antibody

**UnitProt No.**

Q08752

**NCBI Accession No.**

NP\_005029

**Alternative Names**

Peptidyl-prolyl cis-trans isomerase D, Peptidylprolyl isomerase D, 40 kDa peptidyl-prolyl cis-trans isomerase, PPIase D, Cyclophilin-related protein, Rotamase D, CYP-40, CypD, Cyclophilin D

## PRODUCT SPECIFICATION

---

**Antibody Host**

Mouse

**Reacts With**

Human

**Concentration**

1mg/ml (determined by BCA assay)

**Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

**Immunogen**

Recombinant human PPID (1-370aa) purified from E. coli

**Isotype**

IgG2b kappa

**Purification Note**

By protein-A affinity chromatography

**Application**

ELISA, WB, ICC/IF

**Usage**

The antibody has been tested by ELISA, Western blot and ICC/IF analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.

# Human Cyclophilin 40/PPID antibody

Catalog Number: ATGA0473

## Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

## BACKGROUND

### Description


Cyclophilin D (also known as peptidylpropyl isomerase D, PPID) is a member of peptidyl-propyl cis-trans isomerase (PPIase) family, which catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerates the folding of proteins. This protein can bind to the immunosuppressant cyclosporine A and was known that its overexpression suppresses the apoptosis in cancer cells.

### General References

Machida K., et al. (2006) J. Biol. Chem. 281(20): 14314-20.  
Yokoi H., et al. (1996) Genomics. 35(3): 448-55.

## DATA


### Western blot analysis (WB)



The cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human PPID antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: K562 cell lysate  
Lane 2.: Jurkat cell lysate  
Lane 3.: HeLa cell lysate  
Lane 4.: HepG2 cell lysate  
Lane 5.: A549 cell lysate  
Lane 6.: MCF7 cell lysate  
Lane 7.: SK-OV-3 cell lysate  
Lane 8.: PC3 cell lysate

### Immunocytochemistry/Immunofluorescence (ICC/IF)



ICC/IF analysis of PPID in Hep3B cells. The cell was stained with ATGA0473 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).