

# Human CD94/KLRD1 antibody

Catalog Number: ATGA0487

## PRODUCT INFORMATION

---

**Catalog number**

ATGA0487

**Clone No.**

AT13E3

**Product type**

Monoclonal Antibody

**UnitProt No.**

Q13241

**NCBI Accession No.**

NP\_002253

**Alternative Names**

Killer cell lectin like receptor D1, Natural killer cells antigen CD94, KP43, Killer cell lectin-like receptor subfamily D member 1, NK cell receptor

## PRODUCT SPECIFICATION

---

**Antibody Host**

Mouse

**Reacts With**

Human

**Concentration**

1mg/ml (determined by BCA assay)

**Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

**Immunogen**

Recombinant human KLRD1 (32-179aa) purified from E. coli

**Isotype**

IgG1 kappa

**Purification Note**

By protein-A affinity chromatography

**Application**

ELISA, WB, FACS

**Usage**

The antibody has been tested by ELISA, Western blot analysis and Flow cytometry to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.

# Human CD94/KLRD1 antibody

Catalog Number: ATGA0487

## Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

## BACKGROUND

### Description


CD94 (Cluster of Differentiation 94), also known as killer cell lectin-like receptor subfamily D, member 1 (KLRD1) is a human gene. The protein encoded by CD94 gene is a lectin, cluster of differentiation and a receptor that is involved in cell signaling and is expressed on the surface of natural killer cells in the innate immune system. CD94 pairs with the NKG2 molecule as a heterodimer. The CD94/NKG2 complex, on the surface of natural killer cells interacts with Human Leukocyte Antigen (HLA) -E on target cells.

### General References

Cheent KS., et al. (2013) *pnas*. 110(42): 16981-16986.  
Shum BP., et al. (2008) *Nat Immunol*. 168(1): 240-252.  
Fang M., et al. (2011) *Immunity* 34(4): 579-589.


## DATA

### Western blot analysis (WB)



The cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human KLRD1 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: K562 cell lysate  
Lane 2.: MCF7 cell lysate  
Lane 3.: A549 cell lysate




The NK-92 cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human KLRD1 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

### Flow cytometry (FACS)

## Human CD94/KLRD1 antibody

Catalog Number: ATGA0487



Flow cytometry analysis of KLRD1 in PBMC cells. The cell was stained with ATGA0487 at 2-5ug for 1x10<sup>6</sup> cells (red). A Goat anti mouse IgG (Alexa fluor 488) was used as the secondary antibody. Mouse monoclonal IgG was used as the isotype control (dark gray), cells without incubation with primary and secondary antibody was used as the negative control (light gray).