# NKMAXBIO We support you, we believe in your research

# **Human CD34 antibody**

Catalog Number: ATGA0500

# **PRODUCT INFORMATION**

# Catalog number

ATGA0500

#### Clone No.

AT64F9

# **Product type**

Monoclonal Antibody

#### UnitProt No.

P28906

#### **NCBI Accession No.**

NP 001020280

#### **Alternative Names**

Hematopoietic progenitor cell antigen CD34, CD34 molecule

# **PRODUCT SPECIFICATION**

#### **Antibody Host**

Mouse

#### **Reacts With**

Human

#### Concentration

1mg/ml (determined by BCA assay)

#### **Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

#### **Immunogen**

Recombinant human CD34 (32-290aa) purified from E.coli

#### Isotype

IgG2a kappa

#### **Purification Note**

By protein-A affinity chromatography

# **Application**

ELISA, WB, FACS

#### Usage

The antibody has been tested by ELISA, Western blot and FACS analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.

#### **Storage**



# NKMAXBio We support you, we believe in your research

# **Human CD34 antibody**

Catalog Number: ATGA0500

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

### **BACKGROUND**

#### **Description**


Hematopoietic progenitor cell antigen CD34 is a heavily glycosylated, transmembrane glycoprotein that is expressed on the surface of lymphohematopoietic stem and progenitor cells, small-vessel endothelial cells, embryonic fibroblasts and some cells in fetal and adult nervous tissue. It is likely an adhesion molecule with a role in early hematopoiesis by mediating attachment of stem cells to the bone marrow extracellular matrix or directly to stromal cells.

#### **General References**

Satterthwaite A B., et al. (1992) Genomics. 12:788-794. Krause D S., et al. (1994) Blood. 84:691-701.

## **DATA**


## Western blot analysis (WB)



The cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human CD34 antibody (1:500). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: TF1 cell lysate Lane 2.: HeLa cell lysate Lane 3.: Jurkat cell lysate

# Flow cytometry (FACS)



Flow cytometry analysis of CD34 in KG1 cells. The cell was stained with ATGA0500 at 2-5ug for  $1\times10^6$  cells (red). A Goat anti mouse IgG (Alexa fluor 488) was used as the secondary antibody. Mouse monoclonal IgG was used as the isotype control (dark gray), cells without incubation with primary and secondary antibody was used as the negative control (light gray).

