NKMAXBio We support you, we believe in your research

Human NMT1 antibody

Catalog Number: ATGA0501

PRODUCT INFORMATION

Catalog number

ATGA0501

Clone No.

AT2C8

Product type

Monoclonal Antibody

UnitProt No.

P30419

NCBI Accession No.

NP 066565

Alternative Names

Glycylpeptide N-tetradecanoyltransferase 1, Glycylpeptide N-tetradecanoyltransferase 1, NMT, Myristoyl-CoA:protein N-myristoyltransferase 1

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human NMT1 (1-496aa) purified from E. coli

Isotype

IgG2a kappa

Purification Note

By protein-A affinity chromatography

Application

ELISA, WB, ICC/IF

Usage

The antibody has been tested by ELISA, Western blot and ICC/IF analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



NKMAXBio We support you, we believe in your research

Human NMT1 antibody

Catalog Number: ATGA0501

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


Myristate, a rare 14-carbon saturated fatty acid, is cotranslationally attached by an amide linkage to the N-terminal glycine residue of cellular and viral proteins with diverse functions. N-myristoyltransferase catalyzes the transfer of myristate from CoA to proteins. N-myristoylation appears to be irreversible and is required for full expression of the biologic activities of several N-myristoylated proteins, including the alpha subunit of the signal-transducing guanine nucleotide-binding protein (G protein) GO. Recombinant human NMT1 protein, fused to Histag at N-terminus, was expressed in E. coli and purified by using conventional chromatography techniques.

General References

Glover C.J. et al. (1997) J. Biol. Chem. 272:28680-28689. Van Damme P. et al. (2012) Proc. Natl. Acad. Sci. U.S.A. 109:12449-12454.

DATA


Western blot analysis (WB)



The cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human NMT1 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: HepG2 cell lysate Lane 2.: 293T cell lysate

Immunocytochemistry/Immunofluorescence (ICC/IF)



ICC/IF analysis of NMT1 in HeLa cells. The cell was stained with ATGA0501 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

