NKMAXBIO We support you, we believe in your research

Human XPA antibody

Catalog Number: ATGA0502

PRODUCT INFORMATION

Catalog number

ATGA0502

Clone No.

AT71H3

Product type

Monoclonal Antibody

UnitProt No.

P23025

NCBI Accession No.

NP 000371

Alternative Names

DNA repair protein complementing XP-A cells, XP1, XPAC

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human XPA (1-273aa) purified from E. coli

Isotype

IgG1 kappa

Purification Note

By protein-A affinity chromatography

Application

ELISA, WB, ICC/IF

Usage

The antibody has been tested by ELISA, Western blot and ICC/IF analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.

Storage



NKMAXBIO We support you, we believe in your research

Human XPA antibody

Catalog Number: ATGA0502

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

XPA, also known as DNA repair protein complementing XP-A cells, belong to the XPA family. This protein is involved in DNA excision repair. It Initiates repair by binding to damaged sites with various affinities, depending on the photoproduct and the transcriptional state of the region. Defects in XPA are a cause of xeroderma pigmentosum complementation group A (XP-A), which is a rare human autosomal recessive disease characterized by solar sensitivity, high predisposition for developing cancers on areas exposed to sunlight and, in some cases, neurological abnormalities.


General References

Pan Y.R., et al. (2009) Cell Cycle 8:655-664.

James, et al. (2005) Andrews Diseases of the Skin: Clinical Dermatology. (10th ed.). Saunders.

DATA


Western blot analysis (WB)



The cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human XPA antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: SW480 cell lysate Lane 2.: K562 cell lysate Lane 3.: MCF7 cell lysate Lane 4.: LNCaP cell lysate Lane 5.: Raji cell lysate

Immunocytochemistry/Immunofluorescence (ICC/IF)



ICC/IF analysis of XPA in HaLa cells. The cell was stained with ATGA0502 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).

