# NKMAXBIO We support you, we believe in your research

# **Human SETD7 antibody**

Catalog Number: ATGA0524

#### **PRODUCT INFORMATION**

### Catalog number

ATGA0524

#### Clone No.

s4E5

#### **Product type**

Monoclonal Antibody

#### UnitProt No.

Q8WTS6

#### **NCBI Accession No.**

NP 085151

#### **Alternative Names**

SETD7, SET7, SET9, SET7/9, SET7/9 Histone methyltransferase, SET domain-containing protein 8, SET domain-containing protein 7 FLJ21193, SET domain-containing protein 7, Lysine N-methyltransferase 7, Lysine methyltransferase, KMT7, KIAA1717, Histone-lysine N-methyltransferase SETD7, Histone-lysine N-methyltransferase, Histone lysine N methyltransferase H3 lysine 4 specific SET7, Histone lysine methyltransferase, Histone H4-K4 methyltransferase, Histone H3-K4 methyltransferase SETD7, Histone H3 lysine 4 specific methyltransferase, Histone H3 K4 methyltransferase, H4 lysine-4 specific, H3-K4-HMTase SETD7, H3 K4 HMTase, EC 2.1.1.43

#### **Additional Information**

This product was produced from tissue culture supernatant.

#### **PRODUCT SPECIFICATION**

#### **Antibody Host**

Mouse

#### **Reacts With**

Human

#### Concentration

1mg/ml (determined by BCA assay)

#### **Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

#### **Immunogen**

Recombinant human SET7/9 (1-366aa) purified from E. coli

#### Isotype

IgG2b kappa

#### **Purification Note**

By protein-A affinity chromatography



# NKMAXBIO We support you, we believe in your research

## **Human SETD7 antibody**

Catalog Number: ATGA0524

#### **Application**

ELISA, WB, ICC/IF, FACS

#### Usage

The antibody has been tested by ELISA, Western blot, ICC/IF and FACS analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results

#### Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

#### **BACKGROUND**

## **Description**


Set7/9 is a histone methyltransferase (HMTase) that transfers methyl group to Lys4 of histone H3, in complex with S-adenosyl-L-methionine (AdoMet). The methylation of lysine residues of histones plays a critical role in the regulation of chromatin structure and gene expression. Acetylation, phosphorylation and methylation of the amino-terminal tails of histone are thought to be involved in the regulation of chromatin structure and function. The enzymes identified in the methylation of specific lysine residue on histones belong to the SET family with just one exception. Set7/9, unlike most other SET proteins, is exclusively a mono-methylase.

#### **General References**

Bing X, et al., (2003) Nature 421: 652-56. Taewoo K, et al., (2003) EMBO. 22: 292-303. Nishioka K, et al., (2002) Genes Dev. 16: 479-89.

#### **DATA**

#### Western blot analysis (WB)




The Recombinant protein(50ng) was resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human SET7/9 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system. Lane 1.: Recombinant human SET7/9 protein



# **Human SETD7 antibody**

Catalog Number: ATGA0524




The Cell lysates (5ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human SET7/9 antibody (1:2000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: 293T cell lysate


Lane 2.: SET7/9 transfected 293T cell lysate

### Immunocytochemistry/Immunofluorescence (ICC/IF)



ICC/IF analysis of SET7/9 in HeLa cells. The cell was stained with ATGA0524 (1:100). The secondary antibody (green) was used Alexa Fluor 488.

### Flow cytometry (FACS)



Flow cytometry analysis of SET7/9 in Jurkat cells. The cell was stained with ATGA0524 at 2-5ug for 1x10^6cells (red). A Goat anti mouse IgG (Alexa fluor 488) was used as the secondary antibody. Mouse monoclonal IgG was used as the isotype control (darkgray), cells without incubation with primary and secondary antibody was used as the negative control (light gray).

