NKMAXBIO We support you, we believe in your research

Human KAT2A/GCN5 antibody

Catalog Number: ATGA0527

PRODUCT INFORMATION

Catalog number

ATGA0527

Clone No.

AT3G13

Product type

Monoclonal antibody

UnitProt No.

Q92830

NCBI Accession No.

NP 066564

Alternative Names

STAF97, PCAF-b, Lysine acetyltransferase 2A, Histone succinyltransferase KAT2A, Histone glutaryltransferase KAT2A, Histone acetyltransferase GCN5, General control of amino-acid synthesis 5-like 2, General control of amino acid synthesis protein 5-like 2, GCN5L2, GCN5

Additional Information

This product was produced from tissue culture supernatant.

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human GCN5L2 (411-837aa) purified from E. coli

Isotype

IgG2a kappa

Purification Note

By protein-A affinity chromatography

Application

ELISA, WB, ICC/IF



NKMAXBIO We support you, we believe in your research

Human KAT2A/GCN5 antibody

Catalog Number: ATGA0527

Usage

The antibody has been tested by ELISA, Western blot and ICC/IF analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

GCN5L2, as known as GCN5 and KAT2A is the use of immunoprecipitated GCN5L2 for acetylation toprovide the additional components important for acetylation. GCN5L2 can be one of the potential enzymesinvolved in acetylation in vivo. GCN5L2 is expressed predominantly in the embryo and newborn and is essentialfor normal embryonic development. Therefore, GCN5L2 appears to be important for differentiation ofembryonic-derived preadipocytes. On the one hand, loss of GCN5L2 leads to high incidence of apoptosis inthe GCN5L2 mutants that begins before the onset of morphological abnormality.

General References

Wiper-Bergeron N, et al. (2007) Proc Natl Acad Sci U S A, 104(8):2703-8. Jacob AL, et al. (2001) J Biol Chem, 276(40):37659-64. Xu W, et al. (2000) Nat Genet, 26(2):229-232.


DATA

Western blot analysis (WB)




Human KAT2A/GCN5 antibody

Catalog Number: ATGA0527



The cell lysate(35ug) was resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human GCN5L2 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system. Lane 1.: HeLa cell lysate

Immunocytochemistry/Immunofluorescence (ICC/IF)



ICC/IF analysis of GCN5L2 in HeLa cells. The cell was stained with ATGA0527 (1:500). The secondary antibody (green) was used Alexa Fluor 488.

