NKMAXBIO We support you, we believe in your research

Human MafK antibody

Catalog Number: ATGA0538

PRODUCT INFORMATION

Catalog number

ATGA0538

Clone No.

AT2F7

Product type

Monoclonal antibody

UnitProt No.

060675

NCBI Accession No.

NP 002351

Alternative Names

Transcription factor MafK, NFE2u, P18

Additional Information

This product was produced from tissue culture supernatant.

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human MAFK (1-156aa) purified from E. coli

Isotype

IgG2a kappa

Purification Note

By protein-A affinity chromatography

Application

ELISA, WB

Usage

The antibody has been tested by ELISA and Western blot analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



NKMAXBio We support you, we believe in your research

Human MafK antibody

Catalog Number: ATGA0538

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


The developmentally regulated expression of the globin genes depends on upstream regulatory elements termed locuscontrol regions (LCRs). LCRs are associated with powerful enhancer activity that is mediated by thetranscription factor NFE2 (nuclear factor erythroid-2). NFE2 DNA-binding activity consists of aheterodimer containing an 18-kD Maf protein (MafF, MafG or MafK) and p45. Since Maf homodimers lack aputative transactivation domain, the small Mafs behave as transcriptional repressors when they dimerize amongthemselves. However, they seem to serve as transcriptional activators by dimerizing with other (usually larger)basic-zipper proteins and recruiting them to specific DNA-binding sites. Small Maf proteins heterodimerizewith Fos and may act as competitive repressors of the NF-E2 transcription factor. The MafK of Maf proteinsis primarily expressed during development in mesenchymal and hematopoietic cells and neurons.

General References

Oyake, T., et al. (1996) Mol Cell Biol 16(11): 6083-95. Toki, T., et al. (1997) Oncogene 14: 1901-1910.


DATA

Western blot analysis (WB)



The cell lysate(40ug) was resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human MafK antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: Jurkat cell lysate



The recombinant proteins(20ng) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human MafK antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: Recombinant human MafK protein Lane 2.: Recombinant human MafG protein Lane 3.: Recombinant human MafF protein

