NKMAXBIO We support you, we believe in your research

Human IL-33 antibody

Catalog Number: ATGA0555

PRODUCT INFORMATION

Catalog number

ATGA0555

Clone No.

4E9

Product type

Monoclonal antibody

UnitProt No.

095760

NCBI Accession No.

NP 254274

Alternative Names

Interleukin-33, Interleukin-1 family member 11, IL-1F11, Nuclear factor from high endothelial venules, NF-HEV, DVS27-related protein, DVS27, C9orf26

Additional Information

This product was produced from tissue culture supernatant.

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human IL-33 (112-270aa) purified from E. coli

Isotype

IgG2b kappa

Purification Note

By protein-G affinity chromatography

Application

ELISA, WB, FACS

Usage

The antibody has been tested by ELISA, Western blot and FACS analysis to assure specificity and reactivity.



NKMAXBio We support you, we believe in your research

Human IL-33 antibody

Catalog Number: ATGA0555

Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

Interleukin 33 (IL-33) is a 32kDa proinflammatory cytokine and intracellular nuclear factor with transcriptional regulatory properties. IL-33 is structurally related to IL-1, which induces helper T cells to produce type 2 cytokines and acts through the receptor IL1RL-1 (IL1 receptor-like-1), which is known also as ST2. Binding of IL-33 to this receptor activates NF-kappa-B and MAP kinases and induces in vitro Th2 cells to produce cytokines. In vivo, IL-33 induces expression of IL-4, IL-5, IL-13 and leads to severe pathological changes in mucosal organs and in vitro, it can be divided to N-terminal fragment of 12kDa and C-terminal fragment of 18kDa by cleavage of caspase-1.

General References

Brint Ek, et al., (2005) J Biol Chem. 20:277(51). Schmitz, et al., (2005) immunity. 23:479. Baekkevold ES, et al., (2003) Am.J.Pathol. 163(1):69-90.


DATA

Western blot analysis (WB)



Human IL-33 antibody

Catalog Number: ATGA0555




The cell lysate(40ug) was resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human IL-33 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: 293T cell lysate



Human IL-33 antibody


Catalog Number: ATGA0555



The cell lysate(35ug) was resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human IL-33 antibody (1:500). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: Jurkat cell lysate

Flow cytometry (FACS)



Flow cytometry analysis of IL-33 in Jurkat cells. The cell was stained with ATGA0555 at 2-5ug for 1x10^6cells (red). A Goat anti mouse IgG (Alexa fluor 488) was used as the secondary antibody. Mouse monoclonal IgG was used as the isotype control (dark gray), cells without incubation with primary and secondary antibody was used as the negative control (light gray).

