PRODUCT INFORMATION

Catalog number ATGA0569

Clone No. 4D7

Product type Monoclonal antibody

UnitProt No. P19367

NCBI Accession No. NP_000179

Alternative Names

BB404130, Brain form hexokinase, dea, EC 2.7.1.1, Glycolytic enzyme, HEXOKIN, Hexokinase, Hexokinase PI, Hexokinase type I, Hexokinase-A, HK I, HK1 tb, Hk1-s, HK1-ta, HK1-tc, HXK1, mHk1-s, tumor isozyme, Hexokinase 1, Hexokinase 2, Hexokinase 3, Hexokinase 4, HK1,HK2, HK3, HK4

Additional Information

This product was produced from tissue culture supernatant.

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With Human

Concentration 1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human Hexokinase1 (1-917aa) purified from E. coli

Isotype IgG2a kappa

Purification Note By protein-A affinity chromatography

Application ELISA, WB, ICC/IF



Usage

The antibody has been tested by ELISA, Western blot and ICC/IF analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


Hexokinase is the first enzyme in the glycolytic pathway, catalyzing the transfer of a phosphoryl group fromATP to glucose to form glucose-6-phosphate and ADP. In mammals, four distinct enzymes-types 1 to 4hexokinaseshave been identified. The enzyme is found in most cells, but there is tissue specificity for theparticular type of hexokinase. Hexokinase1 is found in the adipose tissue and liver and encodes a ubiquitous formof hexokinase which localizes to the outer membrane of mitochondria. Mutations in this hexokinase1 have beenassociated with hemolytic anemia due to hexokinase deficiency.

General References

Ellison W R. et al., (1975) J Biol Chem 250:1864-1871. Furuta H. et al., (1996) Genomics 36(1):206-9. Jon E. et al., (2003) J.Exp Biology 206:2049-2057.


DATA

Western blot analysis (WB)




The recombinant proteins (20ng) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human Hexokinase antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system. Lane 1.: Recombinant human Hexokinase1 protein

Lane 2.: Recombinant human Hexokinase2 protein Lane 3.: Recombinant human Hexokinase3 protein Lane 4.: Recombinant human Hexokinase4 protein **NKMAXBiO** We support you, we believe in your research **Human Hexokinase antibody** Catalog Number: ATGA0569



The cell lysates(40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human Hexokinase antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system. Lane 1.: HepG2 cell lysate Lane 2.: HeLa cell lysate Lane 3.: Jurkat cell lysate Lane 4.: K562 cell lysate Lane 5.: Mouse brain Tissue lysate

Immunocytochemistry/Immunofluorescence (ICC/IF)



ICC/IF analysis of Hexokinase in HeLa cells. The cell was stained with ATGA0569 (1:100). The secondary antibody (green) was used Alexa Fluor 488. DAPI was stained the cell nucleus (blue).