NKMAXBio We support you, we believe in your research

Human CTLA4 antibody

Catalog Number: ATGA0572

PRODUCT INFORMATION

Catalog number

ATGA0572

Clone No.

AT4C8

Product type

Monoclonal Antibody

UnitProt No.

P16410

NCBI Accession No.

NP 005205

Alternative Names

Cytotoxic T-lymphocyte protein 4 isoform CTLA4-TM precursor, Cytotoxic T-lymphocyte-associated antigen 4, CTLA-4, CD152, CELIAC3, Celiac disease 3, Insulin-dependent diabetes mellitus 12, IDDM12, CD, GSE

PRODUCT SPECIFICATION

Antibody Host

Mouse

Reacts With

Human

Concentration

1mg/ml (determined by BCA assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% glycerol

Immunogen

Recombinant human CTLA4 (36-161aa) purified from E. coli

Isotype

IgG1 kappa

Purification Note

By protein-G affinity chromatography

Application

ELISA, WB

Usage

The antibody has been tested by ELISA and Western blot analysis to assure specificity and reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results.



NKMAXBIO We support you, we believe in your research

Human CTLA4 antibody

Catalog Number: ATGA0572

Storage

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description


CTLA4 (Cytotoxic T-Lymphocyte Antigen 4), also known as CD152, is a protein receptor that downregulates the immune system. CTLA4 is expressed on the surface of Helper T cells and transmits an inhibitory signal to T cells. CTLA4 is similar to the T-cell co-stimulatory protein, CD28, and both molecules bind to CD80 and CD86, also called B7-1 and B7-2 respectively, on antigen-presenting cells. CTLA4 transmits an inhibitory signal to T cells, whereas CD28 transmits a stimulatory signal. Mutations in CTLA4 have been associated with insulin-dependent diabetes mellitus, Graves' disease, Hashimoto's thyroiditis, celiac disease and other autoimmune diseases.

General References

Dariavach. P., et al. (1988) Eur J Immunol 18(12): 1901-1905. Waterhouse. P., et al. (1995) Science 270(5238): 985-988. Magistrelli. G., et al. (1999) Eur J Immunol 29(11): 3596-3602.

DATA

Western blot analysis (WB)



The tissue and cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human CTLA4 antibody (1:500). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.

Lane 1.: Mouse liver tissue lysate

Lane 2.: TF-1 cell lysate Lane 3.: HepG2 cell lysate Lane 4.: WiDr cell lysate

