## NKMAXBio We support you, we believe in your research

# Recombinant human IL-33 protein

Catalog Number: ILC0701

## **PRODUCT INFORMATION**

## **Expression system**

E.coli

#### **Domain**

112-270aa

#### **UniProt No.**

095760

#### **NCBI Accession No.**

NP 254274

#### **Alternative Names**

Interleukin-33, Interleukin-1 family member 11, IL-1F11, Nuclear factor from high endothelial venules, NF-HEV, DVS27-related protein, DVS27, C9orf26

#### **PRODUCT SPECIFICATION**

## **Molecular Weight**

18.1 kDa (160aa) confirmed by MALDI-TOF

#### Concentration

1mg/ml (determined by Bradford assay)

#### **Formulation**

Liquid in. 20mM Tris-HCl buffer (pH 8.0)

#### **Purity**

> 95% by SDS-PAGE

#### **Endotoxin level**

< 1 EU per 1ug of protein (determined by LAL method)

## **Biological Activity**

Measured in a cell proliferation assay using D10.G4.1 mouse helper T cells. The ED50 range ≤ 0.3ng/ml.

#### Tag

Non-Tagged

## **Application**

SDS-PAGE, Bioactivity

#### **Storage Condition**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

## **BACKGROUND**



# NKMAXBio we support you, we believe in your research

## Recombinant human IL-33 protein

Catalog Number: ILC0701

## **Description**

Interleukin 33 (IL-33) is a 32kDa proinflammatory cytokine that may also regulate gene transcription in producer cells. IL-33 is structurally related to IL-1, which induces helper T cells to produce type 2 cytokines and acts through the receptor IL1RL-1 (IL1 receptor-like-1), which is known also as ST2. Binding of IL-33 to this receptor activates NF-kappa-B and MAP kinases and induces in vitro Th2 cells to produce cytokines. In vivo, IL-33 induces expression of IL-4, IL-5, IL-13 and leads to severe pathological changes in mucosal organs and in vitro, it can be divided to N-terminal fragment of 12kDa and C-terminal fragment of 18kDa by cleavage of caspase-1. Recombinant human IL-33(C-terminal fragment) was expressed in E.coli and purified by using conventional chromatography techniques.

#### **Amino acid Sequence**


MSITGISPIT EYLASLSTYN DQSITFALED ESYEIYVEDL KKDEKKDKVL LSYYESQHPS NESGDGVDGK MLMVTLSPTK DFWLHANNKE HSVELHKCEK PLPDQAFFVL HNMHSNCVSF ECKTDPGVFI GVKDNHLALI KVDSSENLCT ENILFKLSET

## **General References**

Schmitz., et al. (2005) immunity 23(5):479-90 Baekkevold, E.S., et al. (2003) Am. J.Pathol. 163(1):69-90

## **DATA**


### **SDS-PAGE**



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

## **Biological Activity**

Human IL-33 (ng/ml)



Human IL-33 in a cell proliferation assay using D10.G4.1 mouse helper T cells. The ED50 range  $\leq$  0.3ng/ml.

