PRODUCT INFORMATION

Expression system E.coli

Domain 134-285aa

UniProt No. Q9Y275

NCBI Accession No. NP_006564.1

Alternative Names

Tumor necrosis factor ligand superfamily member 13B, B lymphocyte stimulator, BLyS, B-cell-activating factor BAFF, Dendritic cell-derived TNF-like molecule, TNF- and APOL-related leukocyte expressed ligand 1, TALL-1, CD257, TNFSF20, ZTNF4

PRODUCT SPECIFICATION

Molecular Weight

21.4 kDa (190aa)

Concentration 1mg/ml (determined by Bradford assay)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4), 10% glycerol

Purity > 95% by SDS-PAGE

Endotoxin level < 1 EU per 1ug of protein (determined by LAL method)

Tag His-Tag

Application SDS-PAGE

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

B lymphocyte stimulator (BLyS), a member of the tumor necrosis factor (TNF) superfamily of cytokines, induces B-cell proliferation and immunoglobulin secretion and is a key regulator of peripheral B-cell populations in vivo..



BLyS is made by immune-cells called monocytes and macrophages. When monocytes and macrophages are activated, BLyS is released and binds to a receptor found only on B cells. Recombinant human BLyS, fused to Histag at N-terminus, was expressed in E. coli and purified by using conventional chromatography techniques.

Amino acid Sequence


<MRGSHHHHHH GMASMTGGQQ MGRDLYDDDD KDRWGSH>MAV QGPEETVTQD CLQLIADSET PTIQKGSYTF VPWLLSFKRG SALEEKENKI LVKETGYFFI YGQVLYTDKT YAMGHLIQRK KVHVFGDELS LVTLFRCIQN MPETLPNNSC YSAGIAKLEE GDELQLAIPR ENAQISLDGD VTFFGALKLL

General References

Allan SM., et al. (2005), Nat. Rev. Immunol. 5:629. Michaud DS., et al. (2006), Cancer Res. 66(8):4525-30

DATA

SDS-PAGE



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

