

Recombinant human GRP58/PDIA3 protein

Catalog Number: ATGP0462

PRODUCT INFORMATION

Expression system

E.coli

Domain

25-505aa

UniProt No.

P30101

NCBI Accession No.

NP_005304.3

Alternative Names

Protein disulfide isomerase family A member 3, Glucose regulated protein 58kDa, GPR58, Protein disulfide isomerase-associated 3, 58 kDa microsomal protein, P58, ERp61, Endoplasmic reticulum resident protein 57, ER protein 57, ERp57, Disulfide isomerase ER-60, Endoplasmic reticulum resident protein 60, ER protein 60, ERp60, GRP57, PI-PLC, HsT17083

PRODUCT SPECIFICATION

Molecular Weight

58.5 kDa (518aa) confirmed by MALDI-TOF

Concentration

1mg/ml (determined by Bradford assay)

Formulation

Liquid in. 20mM Tris-HCl buffer (pH 8.0) containing 1mM DTT, 0.1 M NaCl, and 10% glycerol

Purity

> 95% by SDS-PAGE

Tag

His-Tag

Application

SDS-PAGE

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

PDIA3, also known as protein disulfide-isomerase A3, is a protein of the endoplasmic reticulum that interacts with lectin chaperones calreticulin and calnexin to modulate folding of newly synthesized glycoproteins. This protein has protein disulfide isomerase activity. PDIA3 is also part of the major histocompatibility complex (MHC)

Recombinant human GRP58/PDIA3 protein

Catalog Number: ATGP0462

class I peptide-loading complex (TAP1), which is essential for formation of the final antigen conformation and export from the endoplasmic reticulum to the cell surface. Recombinant human PDIA3 protein, fused to His-tag at N-terminus, was expressed in *E. coli* and purified by using conventional chromatography techniques.

Amino acid Sequence

<MRGSHHHHHH GMASMTGGQQ MGRDLYDDDD KDRWGSM>SDV LELTDDNFES RISDTGSAGL MLVEFFAPWC
GHCKRLAPEY EAAATRLKGI VPLAKVDCTA NTNTCNKYGV SGYPTLKIFR DGEEAGAYDG PRTADGIVSH LKKQAGPASV
PLRTEEEFKK FISDKDASIV GFFDSSFSEA HSEFLKAASN LRDNYRFAHT NVESLVNEYD DNGEGILFR PSHLTNKFED
KTVAYTEQKM TSGKIKKFIQ ENIFGICPHM TEDNKDLIQG KDLLIAYYDV DYEKNAKGSN YWRNRVMMVA KKFLDAGHKL
NFAVASRKTf SHELSDFGLE STAGEIPVVA IRTAKGEKfV MQEEFSRDGK ALERFLQDYF DGNLKRYLKS EPIPESNDGP
VKVVVAENFD EIVNNENKDV LIEFYAPWCG HCKNLEPKYK ELGEKLSKDP NIVIAKMDAT ANDVPSPYEV RGFPTIYFSP
ANKKLNPKKY EGGRELSDFI SYLQREATNP PVIQEEKPKK KKKAQEDL


General References

Vigneron N., et al. (2009) *Eur J Immunol.* 39(9):2371-6.

Forster ML., et al. (2009) *J Biol Chem.* 284(19):13045-56.

DATA

SDS-PAGE



3 μ g by SDS-PAGE under reducing condition and visualized by coomassie blue stain.