NKMAXBIO We support you, we believe in your research

Recombinant human MPG protein

Catalog Number: ATGP1134

PRODUCT INFORMATION

Expression system

E.coli

Domain

1-298aa

UniProt No.

P29372

NCBI Accession No.

NP 002425.2

Alternative Names

DNA-3-methyladenine glycosylase, AAG, ADPG, anpg, APNG, CRA36.1, MDG, Mid1, PIG11, PIG16

PRODUCT SPECIFICATION

Molecular Weight

33.9 kDa (306aa) confirmed by MALDI-TOF (Molecular weight on SDS-PAGE will appear higher)

Concentration

0.25mg/ml (determined by Bradford assay)

Formulation

Liquid in. 20mM Tris-HCl buffer (pH 8.0) containing 1mM DTT, 30% glycerol, 200mM NaCl, 1mM EDTA

Purity

> 90% by SDS-PAGE

Tag

His-Tag

Application

SDS-PAGE

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

MPG (DNA-3-methyladenine glycosylase) belongs to the DNA glycosylase MPG family. This protein initiates base excision repair in DNA by removing a wide variety of alkylated, deaminated, and lipid peroxidation-induced purine adducts. Recombinant human MPG protein, fused to His-tag at C-terminus, was expressed in E. coli and purified by using conventional chromatography techniques.

Amino acid Sequence

MVTPALOMKK PKOFCRRMGO KKORPARAGO PHSSSDAAQA PAEOPHSSSD AAQAPCPRER CLGPPTTPGP YRSIYFSSPK



NKMAXBio We support you, we believe in your research

Recombinant human MPG protein

Catalog Number: ATGP1134


GHLTRLGLEF FDQPAVPLAR AFLGQVLVRR LPNGTELRGR IVETEAYLGP EDEAAHSRGG RQTPRNRGMF MKPGTLYVYI IYGMYFCMNI SSQGDGACVL LRALEPLEGL ETMRQLRSTL RKGTASRVLK DRELCSGPSK LCQALAINKS FDQRDLAQDE AVWLERGPLE PSEPAVVAAA RVGVGHAGEW ARKPLRFYVR GSPWVSVVDR VAEQDTQA<LE HHHHHHH>

General References

Adhikari S., et al. (2009) DNA Repair (Amst). 8(10):1201-6. Allan, J.M., et al. (1998) Cancer Res. 58: 3965-3973.

DATA

SDS-PAGE



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

