PRODUCT INFORMATION

Expression system E.coli

Domain 80-235aa

UniProt No. P06804

NCBI Accession No. NP_038721

Alternative Names

Tumor necrosis factor, Cachectin, TNF-alpha, Tumor necrosis factor ligand superfamily member 2, TNFSF2, TNFa, TNFA, TNF

PRODUCT SPECIFICATION

Molecular Weight

17.4 kDa (157aa) confirmed by MALDI-TOF

Concentration 1mg/ml (determined by Bradford assay)

Formulation Liquid in. Phosphate-Buffered Saline (pH 7.4)

Purity > 95% by SDS-PAGE

Endotoxin level < 1 EU per 1ug of protein (determined by LAL method)

Tag Non-Tagged

Application SDS-PAGE

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

TNFa, also known as tumor necrosis factor, binds to TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. It is mainly secreted by macrophages and can induce cell death of certain tumor cell lines. It is potent pyrogen causing fever by direct action or by stimulation of interleukin-1 secretion and is implicated in the induction of cachexia.



Recombinant mouse TNFa protein was expressed in E. coli and purified by using conventional chromatography.

Amino acid Sequence


MLRSSSQNSS DKPVAHVVAN HQVEEQLEWL SQRANALLAN GMDLKDNQLV VPADGLYLVY SQVLFKGQGC PDYVLLTHTV SRFAISYQEK VNLLSAVKSP CPKDTPEGAE LKPWYEPIYL GGVFQLEKGD QLSAEVNLPK YLDFAESGQV YFGVIAL

General References

Shakhov A.N., et al. (1987) Bioorg. Khim. 13:701-705 Pennica D., et al. (1985) Proc. Natl. Acad. Sci. u.S.A. 82:6060-6064

DATA

SDS-PAGE



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.