# NKMAXBIO We support you, we believe in your research

# Recombinant human ICT1 protein

Catalog Number: ATGP1834

## **PRODUCT INFORMATION**

## **Expression system**

E.coli

#### **Domain**

30-206aa

#### UniProt No.

014197

#### **NCBI Accession No.**

NP 001536

#### **Alternative Names**

Peptidyl-tRNA hydrolase ICT1 mitochondrial, Peptidyl-tRNA hydrolase ICT1, mitochondrial, DS-1, DS1

# **PRODUCT SPECIFICATION**

#### **Molecular Weight**

22.8 kDa (200aa) confirmed by MALDI-TOF

#### Concentration

0.5mg/ml (determined by Bradford assay)

#### **Formulation**

Liquid in. 20mM Tris-HCl buffer (pH 8.0) containing 0.2M NaCl, 30% glycerol, 1mM DTT

#### **Purity**

> 90% by SDS-PAGE

#### Tag

His-Tag

### **Application**

SDS-PAGE

## **Storage Condition**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

#### **BACKGROUND**

#### **Description**

Peptidyl-tRNA hydrolase ICT1 acts as a codon-independent translation release factor that has lost all stop codon specificity and directs the termination of translation in mitochondrion, possibly in case of abortive elongation. It may be involved in the hydrolysis of peptidyl-tRNAs that have been prematurely terminated and thus in the recycling of stalled mitochondrial ribosomes. Recombinant human ICT1 protein, fused to His-tag at N-terminus, was expressed in E. coli and purified by using conventional chromatography techniques.



# NKMAXBio We support you, we believe in your research

# **Recombinant human ICT1 protein**

Catalog Number: ATGP1834

# **Amino acid Sequence**


MGSSHHHHHH SSGLVPRGSH MGSLHKQKDG TEFKSIYSLD KLYPESQGSD TAWRVPNGAK QADSDIPLDR LTISYCRSSG PGGQNVNKVN SKAEVRFHLA TAEWIAEPVR QKIAITHKNK INRLGELILT SESSRYQFRN LADCLQKIRD MITEASQTPK EPTKEDVKLH RIRIENMNRE RLRQKRIHSA VKTSRRVDMD

# **General References**

Richter R., et al. (2010) EMBO J. 29:1116-1125 van Belzen N., et al. (1995) Eur. J. Biochem. 234:843-848

# **DATA**

# **SDS-PAGE**



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

15% SDS-PAGE (3ug)