Recombinant human VEGF R2/KDR/Flk-1 protein

Catalog Number: ATGP3627

PRODUCT INFORMATION

Expression system
Baculovirus

Domain
20-764aa

UniProt No.
P35968

NCBI Accession No.
NP_002244.1

Alternative Names
Vascular endothelial growth factor receptor 2, KDR, CD309, FLK1, VEGFR, VEGFR2

PRODUCT SPECIFICATION

Molecular Weight
110.5 kDa (987aa)

Concentration
0.5mg/ml (determined by absorbance at 280nm)

Formulation
Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

Purity
> 90% by SDS-PAGE

Endotoxin level
< 1 EU per 1ug of protein (determined by LAL method)

Tag
hIgG-His-Tag

Application
SDS-PAGE

Storage Condition
Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description
KDR, also known as vascular endothelial growth factor receptor 2, is one of the subtypes of VEGFR. VEGF receptors are receptors for vascular endothelial growth factor (VEGF). There are three main subtypes of VEGFR (1, 2 and 3). VEGFR2 was shown to be the primary signal transducer for angiogenesis and the development of pathological conditions like cancer. Also, this protein is expressed mainly in the endothelial cells, and the

Recombinant human VEGF R2/KDR/Flk-1 protein

Catalog Number: ATGP3627

expression is upregulated in the tumor vasculature. Recombinant human KDR protein, fused to hIgG-His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

Amino acid Sequence


<ADP>ASVGLPS VSLDLPRLSI QKDILTIKAN TTLQITCRGQ RDLDWLWPNN QSGSEQRVEV TECSDGLFCK TLTIPKVIGN DTGAYKCFYR ETDLASVIYV YVQDYRSPFI ASVSDQHGVV YITENKNKTV VIPCLGSISN LNVSLCARYP EKRFVPDGNR ISWDSKKGFT IPSYMISYAG MVFCEAKIND ESYQSIMYIV VVVGYRIYDV VLSPSHGIEL SVGEKLVLNC TARTELNVGI DFNWEYPSSK HQHKKLVNRD LKTQSGSEMK KFLSTLTIDG VTRSDQGLYT CAASSGLMTK KNSTFVRVHE KPFVAFGSGM ESLVEATVGE RVRIPAKYLG YPPPEIKWYK NGIPLESNHT IKAGHVLTIM EVSERDTGNY TVILTNPISK EKQSHVVSLV VYVPPQIGEK SLISPVDSYQ YGTTQTLTCT VYAIPPPHHI HWYWQLEEEC ANEPSQAVSV TNPYPCEEWR SVEDFQGGNK IEVNKNQFAL IEGKNKTVST LVIQAANVSA LYKCEAVNKV GRGERVISFH VTRGPEITLQ PDMQPTEQES VSLWCTADRS TFENLTWYKL GPQPLPIHVG ELPTPVCKNL DTLWKLNATM FSNSTNDILI MELKNASLQD QGDYVCLAQD RKTKKRHCVV RQLTVLERVA PTITGNLENQ TTSIGESIEV SCTASGNPPP QIMWFKDNET LVEDSGIVLK DGNRNLTIRR VRKEDEGLYT CQACSVLGCA KVEAFFIIEG AQEKTNLE<LE PKSCDKTHTC PPCPAPELLG GPSVFLFPPK PKDTLMISRT PEVTCVVVDV SHEDPEVKFN WYVDGVEVHN AKTKPREEQY NSTYRVVSVL TVLHQDWLNG KEYKCKVSNK ALPAPIEKTI SKAKGQPREP QVYTLPPSRD ELTKNQVSLT CLVKGFYPSD IAVEWESNGQ PENNYKTTPP VLDSDGSFFL YSKLTVDKSR WQQGNVFSCS VMHEALHNHY TQKSLSLSPG KHHHHHH>

General References

Han KY., et al, (2009) Biochem Biophys Res Commun. 382:124-128.
Takyar S., et al, (2016) FASEB J. 30:1317-1327.

DATA

SDS-PAGE



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.