# NKMAXBio We support you, we believe in your research

## Recombinant human ULBP-4/RAET1E protein

Catalog Number: ATGP3656

#### PRODUCT INFORMATION

## **Expression system**

Baculovirus

#### **Domain**

31-225aa

#### UniProt No.

**08TD07** 

#### **NCBI Accession No.**

NP 631904

#### **Alternative Names**

Retinoic acid early transcript 1E, Lymphocyte effector toxicity activation ligand, NKG2D ligand 4, N2DL-4, NKG2DL4, RAE-1-like transcript 4, UL16-binding protein 4, LETAL, N2DL4, ULBP4

## PRODUCT SPECIFICATION

## **Molecular Weight**

49.6 kDa (437aa)

## **Concentration**

0.5mg/ml (determined by absorbance at 280nm)

#### **Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

#### **Purity**

> 85% by SDS-PAGE

#### **Endotoxin level**

< 1 EU per 1ug of protein (determined by LAL method)

## Tag

hlgG-His-Tag

## **Application**

SDS-PAGE

## **Storage Condition**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

## **BACKGROUND**

## Description

RAET1E, also known as NKG2D ligand 4 isoform 1, is a member of the RAET1/ULBP family of cell surface protein that function as ligands for NKG2D. This protein is abnormally expressed on most colon cancer and some other tumor cell lines and virus infected peripheral blood cells. It binds and co-stimulates NKG2D expressing effector



## Recombinant human ULBP-4/RAET1E protein

Catalog Number: ATGP3656

cells including NK cells, NKT cells, gamma delta T cells, and CD8+ alpha beta T cells, activating cytolytic activity and/or cytokine production. Also, it functions as a stress-induced ligand for NKG2D receptor. Recombinant human RAET1E, fused to hlgG-His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.


## **Amino acid Sequence**

ADPHSLCFNF TIKSLSRPGQ PWCEAQVFLN KNLFLQYNSD NNMVKPLGLL GKKVYATSTW GELTQTLGEV GRDLRMLLCD IKPQIKTSDP STLQVEMFCQ REAERCTGAS WQFATNGEKS LLFDAMNMTW TVINHEASKI KETWKKDRGL EKYFRKLSKG DCDHWLREFL GHWEAMPEPT VSPVNASDIH WSSSSLPDVE PKSCDKTHTC PPCPAPELLG GPSVFLFPPK PKDTLMISRT PEVTCVVVDV SHEDPEVKFN WYVDGVEVHN AKTKPREEQY NSTYRVVSVL TVLHQDWLNG KEYKCKVSNK ALPAPIEKTI SKAKGQPREP QVYTLPPSRD ELTKNQVSLT CLVKGFYPSD IAVEWESNGQ PENNYKTTPP VLDSDGSFFL YSKLTVDKSR WQQGNVFSCS VMHEALHNHY TQKSLSLSPG KHHHHHH

#### **General References**

Chalupny NJ., et al, (2003) Biochem. Biophys. Res. Commun. 305:129-135. Kong Y., et al, (2009) Blood 114:310-317.

## **DATA**



15% SDS-PAGE (3ug)

3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

