

Recombinant human Semaphorin 7A protein

Catalog Number: ATGP3720

PRODUCT INFORMATION

Expression system

Baculovirus

Domain

45-648aa

UniProt No.

075326

NCBI Accession No.

NP_003603

Alternative Names

Semaphorin-7A isoform 1, SEMA7A, CD108, CDw108, H-SEMA-K1, H-Sema-L, JMH, SEMAK1, SEMAL

PRODUCT SPECIFICATION

Molecular Weight

95.7 kDa (846aa)

Concentration

0.25mg/ml (determined by absorbance at 280nm)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

Purity

> 90% by SDS-PAGE

Endotoxin level

< 1 EU per 1ug of protein (determined by LAL method)

Tag

hIgG-His-Tag

Application

SDS-PAGE

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

SEMA7A, also known as semaphorin-7A isoform 1, is a membrane-bound semaphorin that associates with cell surfaces via a glycosylphosphatidylinositol (GPI) linkage. It promotes formation of focal adhesion complexes, activation of the protein kinase PTK2/FAK1 and subsequent phosphorylation of MAPK1 and MAPK3. It plays an important role in integrin-mediated signaling and functions both in regulating cell migration and immune

Recombinant human Semaphorin 7A protein

Catalog Number: ATGP3720

responses. It promotes production of proinflammatory cytokines by monocytes and macrophages. Recombinant human SEMA7A, fused to hIgG-His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

Amino acid Sequence

<ADP>QGHLRSG PRIFAVWKGH VGQDRVDFGQ TEPHTVLFHE PGSSSVWVGG RGKVYLFDFP EGKNASVRTV
NIGSTKGSCL DKRDCENYIT LLERRSEGLL ACGTNARHPS CNVLVNGTVV PLGEMRGYAP FSPDENS LVL FEGDEVYSTI
RKQEYNGKIP RFRRIRGESE LYTSDETMQN PQFIKATIVH QDQAYDDKIY YFFREDNPDK NPEAPLNVR VAQLCRGDQG
GESSLVSKW NTFLKAMLVC SDAATNKNFN RLQDVFLLPD PSGQWRDTRV YGVFSNPWNY SAVCVYSLGD IDKVFRTSSL
KGYHSSLPNP RPGKCLPDQQ PIPTETFQVA DRHPEVAQRV EPMGPLKTPL FHSKYHYQKV AVHRMQASHG ETFHVLYLT
DRGTIHVKVE PGEQEHSFAF NIMEIQPFRR AAAIQTMSLD AERRKLYVSS QWEVSQVPLD LCEVYGGGCH GCLMSRDPYC
GWDQGRCISI YSSERSVQLQS INPAEPHKEC PNPKPDKAPL QKVSLAPNSR YYLSCPME SR HATYSWRHKE NVEQSCEPGH
QSPNCILFIE NLTAQQYGHY FCEAQEGSYF REAQHWQLLP EDGIMAEHLL GHACALA<LEP KSCDKHTCP PCPAPELLGG
PSVFLFPPKP KDTLMISRTP EVTCVVVDVS HEDPEVKFNW YVDGVEVHNA KTKPREEQYN STYRVVSVLT VLHQDWLNGK
EYKCKVSNKA LPAPIEK TIS KAKGQP REPQ VYTLPPSRDE LTKNQVSLTC LVKGFYPSDI AVEWESNGQP ENNYKTTPPV
LSDGSFFLY SKLTVDKSRW QQGNVFSCSV MHEALHNHYT QKSLSLSPGK HHHHHH>


General References

Scott GA., et al. (2008) J Invest Dermatol. 128:151-161.

Pasterkamp RJ., et al. (2003) Nature. 424:398-405.

DATA

SDS-PAGE



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain