NKMAXBio We support you, we believe in your research

Recombinant human IL-18R alpha/IL18R1 protein

Catalog Number: ATGP3821

PRODUCT INFORMATION

Expression system

Baculovirus

Domain

19-329aa

UniProt No.

013478

NCBI Accession No.

NP 003846

Alternative Names

Interleukin-18 receptor 1, CD218 antigen-like family member A, CDw218a, IL1 receptor-related protein, IL-1Rrp, IL1R-rp, Interleukin-18 receptor alpha, IL-18R-alpha, IL-18Ralpha, CD218a, IL1RRP, IL-1Rrp

PRODUCT SPECIFICATION

Molecular Weight

62.7 kDa (550aa)

Concentration

0.5mg/ml (determined by absorbance at 280nm)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

Purity

> 90% by SDS-PAGE

Endotoxin level

< 0.1 EU per lug of protein (determined by LAL method)

Tag

hlgG-His-Tag

Application

SDS-PAGE

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

IL18R1, also known as interleukin 18 receptor 1, is an interleukin receptor of the immunoglobulin superfamily. This protein contain three Ig-like C2-type (immunoglobulin-like) domains and on TIR domain. It shares numerous immuno-regulatory functions with IL12. IFN-alpha and IL12 are reported to induce the expression of this receptor



NKMAXBIO We support you, we believe in your research

Recombinant human IL-18R alpha/IL18R1 protein

Catalog Number: ATGP3821

in NK and T cells. Also, this receptor specifically binds interleukin 18 (IL18), and is essential for IL18 mediated signal transduction. It binds to the agonist leads to the activation of NF-kappa-B, but does not bind IL1A or IL1B beta. Recombinant human IL18R1, fused to hIgG-His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

Amino acid Sequence


AESCTSRPHI TVVEGEPFYL KHCSCSLAHE IETTTKSWYK SSGSQEHVEL NPRSSSRIAL HDCVLEFWPV ELNDTGSYFF QMKNYTQKWK LNVIRRNKHS CFTERQVTSK IVEVKKFFQI TCENSYYQTL VNSTSLYKNC KKLLLENNKN PTIKKNAEFE DQGYYSCVHF LHHNGKLFNI TKTFNITIVE DRSNIVPVLL GPKLNHVAVE LGKNVRLNCS ALLNEEDVIY WMFGEENGSD PNIHEEKEMR IMTPEGKWHA SKVLRIENIG ESNLNVLYNC TVASTGGTDT KSFILVRKAD MADIPGHVFT RVEPKSCDKT HTCPPCPAPE LLGGPSVFLF PPKPKDTLMI SRTPEVTCVV VDVSHEDPEV KFNWYVDGVE VHNAKTKPRE EQYNSTYRVV SVLTVLHQDW LNGKEYKCKV SNKALPAPIE KTISKAKGQP REPQVYTLPP SRDELTKNQV SLTCLVKGFY PSDIAVEWES NGQPENNYKT TPPVLDSDGS FFLYSKLTVD KSRWQQGNVF SCSVMHEALH NHYTQKSLSL SPGKHHHHHH

General References

Torigoe K., et al, (1997) J. Biol. Chem. 272:25737-25742. Hoshino K., et al, (1999) J. Immunol. 162:5041-5044.

DATA

SDS-PAGE



15% SDS-PAGE (3ug)

3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

