PRODUCT INFORMATION

Expression system Baculovirus

Domain 34-179aa

UniProt No. Q9GZX6

NCBI Accession No. NP_065386

Alternative Names

Interleukin-22, IL22, IL-21, IL-22, IL-D110, IL-TIF, ILTIF, TIFa, TIFIL-23, zcyto18, Macrophagleukin-22, IL22, 2

PRODUCT SPECIFICATION

Molecular Weight 17.8 kDa (155aa)

Concentration 1mg/ml (determined by absorbance at 280nm)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

Purity

> 90% by SDS-PAGE

Endotoxin level < 1 EU per 1ug of protein (determined by LAL method)

Biological Activity

The activity is determined by the IL-10 ELISA in a using COLO 205 human colorectal adenocarcinoma cells. The ED50 range \leq 1.2ng/ml.

Tag His-Tag

Application SDS-PAGE, Bioactivity

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

IL22, also known as interleukin-22, is a member of a group of cytokines involved the IL-10 family or IL-10 subfamily. This protein shares us of IL-10R2 in cell signaling with other members of this family, IL-10, IL-26, IL-28A/B and IL-29. It is produced by normal T cells upon anti-CD3 stimulation and initiates innate immune responses against bacterial pathogens especially in epithelial cells such as respiratory and gut epithelial cells. Also, it can be produced by Th17 cells and play a role in the coordinated response of both adaptive and innate immune systems. Biological activity of IL-22 is the functional receptor complex consists of two receptor subunits, IL-22R and IL-10R and is further regulated by interactions with a soluble binding protein, IL-22BP, which shares sequence similarity with an extracellular region of IL-22R1. Recombinant human IL22, fused to His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

Amino acid Sequence


<ADP>APISSHC RLDKSNFQQP YITNRTFMLA KEASLADNNT DVRLIGEKLF HGVSMSERCY LMKQVLNFTL EEVLFPQSDR FQPYMQEVVP FLARLSNRLS TCHIEGDDLH IQRNVQKLKD TVKKLGESGE IKAIGELDLL FMSLRNACI<H HHHHH>

General References

Kontenko SV., et al, (2001) J. Biol. Chem. 276:2725-2732. Xie MH., et al, (2000) J. Biol. Chem. 275:31335-31339.

DATA

SDS-PAGE



Biological Activity

3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

NKMAXBiO We support you, we believe in your research Recombinant human IL-22 protein Catalog Number: ATGP3842

Human IL-22 (ng/ml) 2.5 1.5 0.5 0.6 0.6 10^4 10^3 10^2 10^1 10^9 10^1 10^2 10^3 10^4 10^5

The activity is determined by the IL-10 ELISA in a using COLO 205 human colorectal adenocarcinoma cells. The ED50 range \leq 1.2 ng/ml.