# NKMAXBIO We support you, we believe in your research

# **Recombinant mouse CD200 protein**

Catalog Number: ATGP3878

#### PRODUCT INFORMATION

## **Expression system**

Baculovirus

#### **Domain**

31-232aa

#### **UniProt No.**

054901

#### **NCBI Accession No.**

NP 034948

#### **Alternative Names**

Cd200, Mox2, OX2, OX-2 membrane glycoprotein isoform 1, MRC OX-2 antigen

# **PRODUCT SPECIFICATION**

#### **Molecular Weight**

49.5 kDa (441aa)

#### Concentration

1mg/ml (determined by absorbance at 280nm)

#### **Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

#### **Purity**

> 95% by SDS-PAGE

#### **Endotoxin level**

< 1 EU per 1ug of protein (determined by LAL method)

#### Tag

hlgG-His-Tag

## **Application**

SDS-PAGE

# **Storage Condition**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

#### **BACKGROUND**

#### **Description**

CD200, also known as OX-2 membrane glycoprotein isoform 1, is one of the immunoglobulin superfamily. It plays an important role in anti-tumor immunity through interaction with its receptor, and overexpression of this protein effects on malignancies as well as on cancer stem cells. In T cells, this protein functions as a costimulatory effect that is independent of the CD28 pathway. It also plays an important role on myeloid cell



# NKMAXBio we support you, we believe in your research

# **Recombinant mouse CD200 protein**

Catalog Number: ATGP3878

regulation that is suggested by restricted expression of its receptor. Studies of this protein in mouse and rat suggest that this gene may regulate myeloid cell activity and delivers an inhibitory signal for the macrophage lineage in diverse tissues. Recombinant mouse CD200, fused to hlgG-His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

## **Amino acid Sequence**


QVEVVTQDER KALHTTASLR CSLKTSQEPL IVTWQKKKAV SPENMVTYSK THGVVIQPAY KDRINVTELG LWNSSITFWN TTLEDEGCYM CLFNTFGSQK VSGTACLTLY VQPIVHLHYN YFEDHLNITC SATARPAPAI SWKGTGTGIE NSTESHFHSN GTTSVTSILR VKDPKTQVGK EVICQVLYLG NVIDYKQSLD KG<LEPKSCDK THTCPPCPAP ELLGGPSVFL FPPKPKDTLM> <ISRTPEVTCV VVDVSHEDPE VKFNWYVDGV EVHNAKTKPR EEQYNSTYRV VSVLTVLHQD> <WLNGKEYKCK VSNKALPAPI EKTISKAKGQ PREPQVYTLP PSRDELTKNQ VSLTCLVKGF> <YPSDIAVEWE SNGQPENNYK TTPPVLDSDG SFFLYSKLTV DKSRWQQGNV FSCSVMHEAL> <HNHYTQKSLS LSPGKHHHHH H>

#### **General References**

Poongodi R., et al. (2018) Indian J Pathol Microbiol. 61:50-57. Kuwabara J., et al. (2018) Biochem Biophys Res Commun. 496:542-548.

## **DATA**

#### **SDS-PAGE**



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

