

Recombinant human CTLA4 protein

Catalog Number: ATGP3914

PRODUCT INFORMATION

Expression system

Baculovirus

Domain

36-161aa

UniProt No.

P16410

NCBI Accession No.

NP_005205

Alternative Names

Cytotoxic T-lymphocyte protein 4 isoform CTLA4-TM precursor, Cytotoxic T-lymphocyte-associated antigen 4, CTLA-4, CD152, CELIAC3, Celiac disease 3, Insulin-dependent diabetes mellitus 12, IDDM12, CD, GSE

Additional Information

ATGP3364 has been replaced with a catalog number ATGP3914.

PRODUCT SPECIFICATION

Molecular Weight

40.8 kDa (368aa)

Concentration

0.5mg/ml (determined by absorbance at 280nm)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

Purity

> 95% by SDS-PAGE

Endotoxin level

< 1 EU per 1ug of protein (determined by LAL method)

Biological Activity

The activity is determined by the IL-2 ELISA in a using Jurkat human acute T cell leukemia cells. The ED50 range \leq 150ng/ml with Human B7 1/CD80.

Tag

hIgG-His-Tag

Application

SDS-PAGE, Bioactivity

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

Recombinant human CTLA4 protein

Catalog Number: ATGP3914

BACKGROUND

Description

CTLA-4, also known as cytotoxic T-lymphocyte protein 4, is a type I transmembrane T cell inhibitory molecule that is a member of the Ig superfamily. The affinity of CTLA-4 for its natural B7 family ligands, CD80 and CD86, is considerably stronger than the affinity of their cognate stimulatory coreceptor CD28. Recombinant human CTLA-4, fused to hIgG His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

Amino acid Sequence


```
<ADL>KAMHVAQ PAVVLASSRG IASFVCEYAS PGKATEVRVT VLRQADSQVT EVCAATYMMG NELTFLDDSI
CTGTSSGNQV NLTIQGLRAM DTGLYICKVE LMYPPTYLG IGNGTQIYVI DPEPCPDS<L EPKSCDKTHT CPPCPAPELL
GGPSVFLFPP KPKDTLMISR TPEVTCVVVD VSHEDPEVKF NWYVDGVEVH NAKTKPREEQ YNSTYRVVSV LTVLHQDWLN
GKEYKCKVSN KALPAIEKT ISKAKGQPRE PQVYTLPPSR DELTKNQVSL TCLVKGFYPS DIAVEWESNG QPENNYKTP
PVLDSGDSFF LYSKLTVDKS RWQQGNVFSC SVMHEALHNNH YTQKSLSLSP GKHHHHHHH>
```

General References

Harper K., et al. (1991) J. Immunol. 147:1037-1044.
 Teft WA., et al. (2006) Annu. Rev. Immunol. 24:65-97.


DATA

SDS-PAGE



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

Biological Activity



Human CTLA-4 inhibits Human B7 1/CD80 induced IL-2 secretion of the Jurkat human acute T cell leukemia cells. The ED50 range ≤150 ng/ml.