# NKMAXBIO We support you, we believe in your research

# Recombinant human CD200 protein

Catalog Number: ATGP4013

#### PRODUCT INFORMATION

### **Expression system**

Baculovirus

#### **Domain**

31-232aa

#### UniProt No.

P41217

#### **NCBI Accession No.**

NP 005935

#### **Alternative Names**

OX-2 membrane glycoprotein isoform a, CD200, MOX1, MOX2, MRC, OX-2

#### PRODUCT SPECIFICATION

#### **Molecular Weight**

23.5 kDa (211aa)

#### Concentration

0.5mg/ml (determined by Absorbance at 280nm)

#### **Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

#### **Purity**

> 95% by SDS-PAGE

#### **Endotoxin level**

< 1 EU per 1ug of protein (determined by LAL method)

#### **Biological Activity**

Measured by its binding ability in a functional ELISA with Human CD200R1 (CAT# ATGP3809)

#### Tag

His-Tag

#### **Application**

SDS-PAGE, Bioactivity

### **Storage Condition**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

# **BACKGROUND**

# Description

CD200, also known as OX-2 membrane glycoprotein isoform a, belongs to the immunoglobulin superfamily. This



# Recombinant human CD200 protein

Catalog Number: ATGP4013

protein imparts immune privileges by suppressing alloimmune and autoimmune responses through its receptor, CD200R, expressed primarily on myeloid cells. D200 and CD200R associate via their respective N-terminal Ig-like domains. CD200-CD200R signaling have been shown to play an important role in the regulation of anti-tumor immunity, and overexpression of CD200 has been reported in a number of malignancies, including CLL, as well as on cancer stem cells. Studies of the related genes in mouse and rat suggest that this gene may regulate myeloid cell activity and delivers an inhibitory signal for the macrophage lineage in diverse tissues. Elevated expression of this protein has been reported to be associated with poor prognosis in a number of human malignancies. In addition, it also plays an important role in prevention of graft rejection, autoimmune diseases and spontaneous abortion. Recombinant human CD200 protein, fused to His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

#### **Amino acid Sequence**


<ADP>QVQVVTQ DEREQLYTPA SLKCSLQNAQ EALIVTWQKK KAVSPENMVT FSENHGVVIQ PAYKDKINIT QLGLQNSTIT FWNITLEDEG CYMCLFNTFG FGKISGTACL TVYVQPIVSL HYKFSEDHLN ITCSATARPA PMVFWKVPRS GIENSTVTLS HPNGTTSVTS ILHIKDPKNQ VGKEVICQVL HLGTVTDFKQ TVNKG

#### **General References**

Akkaya M., et al. (2013) PLoS One. 8:e63325. Li L., et al. (2016) Med Sci Monit. 22:1079-1084.

#### **DATA**


#### **SDS-PAGE**



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

## **Biological Activity**

Human CD200R1 (ug/ml)



CD200 Subunit is coated at 10 ug/ml (100 ul/well) can bind Human CD200R1 (CAT# ATGP3809) in a Functional ELISA assay.

